

MINKOWSKI TYPE PROBLEMS FOR CONVEX HYPERSURFACES IN HYPERBOLIC SPACE

CLAUS GERHARDT

ABSTRACT. We consider the problem $F = f(\nu)$ for strictly convex, closed hypersurfaces in \mathbb{H}^{n+1} and solve it for curvature functions F the inverses of which are of class (K^*) .

CONTENTS

0. Introduction	1
1. Notations and definitions	3
2. The Beltrami map	6
3. Hadamard's theorem in hyperbolic space	10
4. The Gauß maps	12
5. Curvature flow	21
6. Curvature estimates	28
7. Convergence to a stationary solution	30
References	31

0. INTRODUCTION

In the classical Minkowski problem in \mathbb{R}^{n+1} one wants to find a strictly convex closed hypersurface $M \subset \mathbb{R}^{n+1}$ such that its Gauß curvature K equals a given function f defined in the normal space of M or equivalently defined on S^n

$$(0.1) \quad K|_M = f(\nu).$$

The problem has been partially solved by Minkowski [12], Alexandrov [1], Lewy [11], Nirenberg [13], and Pogorelov [15], and in full generality by Cheng and Yau [2].

Instead of prescribing the Gaussian curvature other curvature functions F can be considered, i.e., one studies the problem

$$(0.2) \quad F|_M = f(\nu).$$

Date: April 6, 2006.

2000 Mathematics Subject Classification. 35J60, 53C21, 53C44, 53C50, 58J05.

Key words and phrases. de Sitter space-time, hyperbolic space, Minkowski problem.

If F is one of the symmetric polynomials H_k , $1 \leq k \leq n$, this problem has been solved by Guan and Guan [9]. They proved that (0.2) has a solution, if f is invariant with respect to a fixed point free group of isometries of S^n .

In a previous work [8] we solved the problem (0.2) for strictly convex hypersurfaces $M \subset S^{n+1}$ and for curvature functions F the inverses of which are of class (K) , see [6, Definition 1.3]. These F include all H_k , $1 \leq k \leq n$, $|A|^2$, and also any symmetric, convex curvature function homogeneous of degree 1, cf. [5, Lemma 1.6].

In the present paper we consider the problem (0.2) for strictly convex hypersurfaces in \mathbb{H}^{n+1} and for curvature functions F the inverses of which belong to a subclass of (K) , the so-called class (K^*) , cf. [6, Definition 1.6].

Among the curvature functions F that satisfy this requirement are the Gauss curvature $F = K = H_n$, and all curvature functions that can be written as

$$(0.3) \quad F = H_k K^a, \quad 1 \leq k \leq n,$$

where $a > 0$ is a constant, as well as positive powers of those functions.

The Minkowski space $\mathbb{R}^{n+1,1}$ contains two spaces of constant curvature as hypersurfaces, namely, \mathbb{H}^{n+1} which is defined as

$$(0.4) \quad \mathbb{H}^{n+1} = \{x \in \mathbb{R}^{n+1,1} : \langle x, x \rangle = -1, x^0 > 0\}$$

and the de Sitter space-time N , a Lorentzian manifold of constant curvature $K_N = 1$

$$(0.5) \quad N = \{x \in \mathbb{R}^{n+1,1} : \langle x, x \rangle = 1\}.$$

We shall show in Section 4 that for any closed strictly convex hypersurface $M \subset \mathbb{H}^{n+1}$ there exists a Gauß map

$$(0.6) \quad x \in M \rightarrow \tilde{x} \in M^* \subset N,$$

where M^* is the polar set of M . M^* is spacelike, also strictly convex, as smooth as M , and the Gauß map is a diffeomorphism.

On the other hand, for any given closed, spacelike, connected, strictly convex hypersurface $M \subset N$ there also exists a Gauß map

$$(0.7) \quad x \in M \rightarrow \tilde{x} \in M^* \subset \mathbb{H}^{n+1}$$

which maps M onto a closed, strictly convex hypersurface in hyperbolic space. These Gauß maps are inverse to each other.

If we consider $M \subset \mathbb{H}^{n+1}$ as an embedding in $\mathbb{R}^{n+1,1}$ of codimension 2, so that the tangent spaces $T_x(M)$ and $T_x(\mathbb{H}^{n+1})$ can be identified with subspaces of $T_x(\mathbb{R}^{n+1,1})$, then the image of the point x under the Gauß map is exactly the normal vector $\nu \in T_x(\mathbb{H}^{n+1})$

$$(0.8) \quad \tilde{x} = \nu \in T_x(\mathbb{H}^{n+1}) \subset T_x(\mathbb{R}^{n+1,1}).$$

Thus, the equation (0.2) can also be written in the form

$$(0.9) \quad F|_M = f(\tilde{x}) \quad \forall x \in M,$$

where f is given as a function defined in N .

Using (0.6) we shall prove that (0.9) has a dual problem, namely,

$$(0.10) \quad \tilde{F}|_{M^*} = f^{-1}(\tilde{x}) \quad \forall \tilde{x} \in M^*,$$

where \tilde{F} is the inverse of F

$$(0.11) \quad \tilde{F}(\kappa_i) = \frac{1}{F(\kappa_i^{-1})}.$$

In the dual problem the curvature is not prescribed by a function defined in the normal space, but by a function defined on the hypersurface.

Both problems are equivalent, solving one also leads to a solution of the dual one; notice also that

$$(0.12) \quad M^{**} = M \quad \wedge \quad \tilde{\tilde{x}} = x.$$

To find a solution we assume that (\tilde{F}, f^{-1}) satisfy barrier conditions, cf. Definition 5.3 for details.

Then we shall prove

0.1. Theorem. *Let $F \in C^{m,\alpha}(\Gamma_+)$, $2 \leq m$, $0 < \alpha < 1$, be a symmetric, positively homogeneous and monotone curvature function such that its inverse \tilde{F} is of class (K^*) , let $0 < f \in C^{m,\alpha}(N)$ and assume that the barrier conditions for (\tilde{F}, f^{-1}) are satisfied, then the dual problems*

$$(0.13) \quad F|_M = f(\tilde{x})$$

and

$$(0.14) \quad \tilde{F}|_{M^*} = f^{-1}(\tilde{x})$$

have strictly convex solutions M resp. M^* of class $C^{m+2,\alpha}$, where M^* is spacelike.

The paper is organized as follows: Section 1 gives an overview of the definitions and conventions we rely on. In Section 2 we define the Beltrami map from \mathbb{H}^{n+1} to \mathbb{R}^{n+1} with the help of which we prove Hadamard's theorem for strictly convex hypersurfaces in \mathbb{H}^{n+1} in Section 3.

The Gauß maps and their properties are treated in Section 4. In the last three sections we prove the existence of a solution in N using a curvature flow method.

1. NOTATIONS AND DEFINITIONS

The main objective of this section is to state the equations of Gauß, Codazzi, and Weingarten for hypersurfaces. Since we are dealing with hypersurfaces in a Riemannian space as well as in a Lorentzian space, we shall formulate the governing equations of a hypersurface M in a semi-riemannian $(n+1)$ -dimensional manifold N , which is either Riemannian or Lorentzian. Geometric quantities in N will be denoted by $(\bar{g}_{\alpha\beta}), (\bar{R}_{\alpha\beta\gamma\delta})$, etc., and those in M by $(g_{ij}), (R_{ijkl})$, etc. Greek indices range from 0 to n and Latin from 1 to n ; the summation convention is always used. Generic coordinate systems

in N resp. M will be denoted by (x^α) resp. (ξ^i) . Covariant differentiation will simply be indicated by indices, only in case of possible ambiguity they will be preceded by a semicolon, i.e., for a function u in N , (u_α) will be the gradient and $(u_{\alpha\beta})$ the Hessian, but e.g., the covariant derivative of the curvature tensor will be abbreviated by $\bar{R}_{\alpha\beta\gamma\delta;\epsilon}$. We also point out that

$$(1.1) \quad \bar{R}_{\alpha\beta\gamma\delta;i} = \bar{R}_{\alpha\beta\gamma\delta;\epsilon} x_i^\epsilon$$

with obvious generalizations to other quantities.

Let M be a *spacelike* hypersurface, i.e. the induced metric is Riemannian, with a differentiable normal ν . We define the signature of ν , $\sigma = \sigma(\nu)$, by

$$(1.2) \quad \sigma = \bar{g}_{\alpha\beta} \nu^\alpha \nu^\beta = \langle \nu, \nu \rangle.$$

In case N is Lorentzian, $\sigma = -1$, and ν is time-like.

In local coordinates, (x^α) and (ξ^i) , the geometric quantities of the spacelike hypersurface M are connected through the following equations

$$(1.3) \quad x_{ij}^\alpha = -\sigma h_{ij} \nu^\alpha$$

the so-called *Gauß formula*. Here, and also in the sequel, a covariant derivative is always a *full* tensor, i.e.

$$(1.4) \quad x_{ij}^\alpha = x_{,ij}^\alpha - \Gamma_{ij}^k x_k^\alpha + \bar{\Gamma}_{\beta\gamma}^\alpha x_i^\beta x_j^\gamma.$$

The comma indicates ordinary partial derivatives.

In this implicit definition the *second fundamental form* (h_{ij}) is taken with respect to $-\sigma\nu$.

The second equation is the *Weingarten equation*

$$(1.5) \quad \nu_i^\alpha = h_i^k x_k^\alpha,$$

where we remember that ν_i^α is a full tensor.

Finally, we have the *Codazzi equation*

$$(1.6) \quad h_{ij;k} - h_{ik;j} = \bar{R}_{\alpha\beta\gamma\delta} \nu^\alpha x_i^\beta x_j^\gamma x_k^\delta$$

and the *Gauß equation*

$$(1.7) \quad R_{ijkl} = \sigma \{h_{ik} h_{jl} - h_{il} h_{jk}\} + \bar{R}_{\alpha\beta\gamma\delta} x_i^\alpha x_j^\beta x_k^\gamma x_l^\delta.$$

Here, the signature of ν comes into play.

Now, let us assume that N is a globally hyperbolic Lorentzian manifold with a *compact* Cauchy surface. Then N is a topological product $\mathbb{R} \times \mathcal{S}_0$, where \mathcal{S}_0 is a compact Riemannian manifold, and there exists a Gaussian coordinate system (x^α) , such that the metric in N has the form

$$(1.8) \quad d\bar{s}_N^2 = e^{2\psi} \{-dx^{0^2} + \sigma_{ij}(x^0, x) dx^i dx^j\},$$

where σ_{ij} is a Riemannian metric, ψ a function on N , and x an abbreviation for the spacelike components (x^i) ,

We also assume that the coordinate system is *future oriented*, i.e. the time coordinate x^0 increases on future directed curves. Hence, the *contravariant* time-like vector $(\xi^\alpha) = (1, 0, \dots, 0)$ is future directed as is its *covariant* version $(\xi_\alpha) = e^{2\psi}(-1, 0, \dots, 0)$.

Let $M = \text{graph } u|_{\mathcal{S}_0}$ be a spacelike hypersurface

$$(1.9) \quad M = \{ (x^0, x) : x^0 = u(x), x \in \mathcal{S}_0 \},$$

then the induced metric has the form

$$(1.10) \quad g_{ij} = e^{2\psi} \{-u_i u_j + \sigma_{ij}\}$$

where σ_{ij} is evaluated at (u, x) , and its inverse $(g^{ij}) = (g_{ij})^{-1}$ can be expressed as

$$(1.11) \quad g^{ij} = e^{-2\psi} \left\{ \sigma^{ij} + \frac{u^i u^j}{v} \right\},$$

where $(\sigma^{ij}) = (\sigma_{ij})^{-1}$ and

$$(1.12) \quad \begin{aligned} u^i &= \sigma^{ij} u_j \\ v^2 &= 1 - \sigma^{ij} u_i u_j \equiv 1 - |Du|^2. \end{aligned}$$

Hence, graph u is spacelike if and only if $|Du| < 1$.

The covariant form of a normal vector of a graph looks like

$$(1.13) \quad (\nu_\alpha) = \pm v^{-1} e^\psi (1, -u_i).$$

and the contravariant version is

$$(1.14) \quad (\nu^\alpha) = \mp v^{-1} e^{-\psi} (1, u^i).$$

Thus, we have

1.1. Remark. Let M be spacelike graph in a future oriented coordinate system. Then, the contravariant future directed normal vector has the form

$$(1.15) \quad (\nu^\alpha) = v^{-1} e^{-\psi} (1, u^i)$$

and the past directed

$$(1.16) \quad (\nu^\alpha) = -v^{-1} e^{-\psi} (1, u^i).$$

In the Gauß formula (1.3) we are free to choose the future or past directed normal, but we stipulate that in general we use the past directed normal unless otherwise stated.

Look at the component $\alpha = 0$ in (1.3), then we obtain in view of (1.16)

$$(1.17) \quad e^{-\psi} v^{-1} h_{ij} = -u_{ij} - \bar{I}_{00}^0 u_i u_j - \bar{I}_{0i}^0 u_j - \bar{I}_{0j}^0 u_i - \bar{I}_{ij}^0.$$

Here, the covariant derivatives are taken with respect to the induced metric of M , and

$$(1.18) \quad -\bar{I}_{ij}^0 = e^{-\psi} \bar{h}_{ij},$$

where (\bar{h}_{ij}) is the second fundamental form of the hypersurfaces $\{x^0 = \text{const}\}$.

2. THE BELTRAMI MAP

Let $\mathbb{R}^{n+1,1}$ be the $(n+2)$ -dimensional Minkowski space with points $x = (x^a)$, $0 \leq a \leq n+1$, where x^0 is the time function.

The submanifolds

$$(2.1) \quad \mathbb{H}^{n+1} = \{x \in \mathbb{R}^{n+1,1} : \langle x, x \rangle = -1, x^0 > 0\}$$

and

$$(2.2) \quad N = \{x \in \mathbb{R}^{n+1,1} : \langle x, x \rangle = 1\}$$

are spaces of constant curvature. \mathbb{H}^{n+1} is the $(n+1)$ -dimensional hyperbolic space with constant curvature $K = -1$, and N is a Lorentzian manifold with constant curvature $K_N = 1$, the de Sitter space-time.

N is globally hyperbolic, as can be seen by introducing polar coordinates in the Euclidean part of the Minkowski space such that the metric in $\mathbb{R}^{n+1,1}$ is expressed as

$$(2.3) \quad d\bar{s}^2 = -dx^{02} + dr^2 + r^2 \sigma_{ij} d\xi^i d\xi^j,$$

where σ_{ij} is the metric in S^n .

Then N is the embedding

$$(2.4) \quad N = \{(x^0, r, \xi^i) : r = \sqrt{1 + |x^0|^2}, x^0 \in \mathbb{R}, \xi \in S^n\},$$

i.e., $N = \mathbb{R} \times S^n$ topologically and

$$(2.5) \quad ds_N^2 = e^{2\psi} \{-d\tau^2 + \sigma_{ij} d\xi^i d\xi^j\},$$

where

$$(2.6) \quad \tau = \int_0^{x^0} \frac{1}{1+t^2} \quad \wedge \quad \psi = \frac{1}{2} \log(1 + |x^0|^2).$$

Notice that N is simply connected, since $n \geq 2$.

Let us analyze a special representation of \mathbb{H}^{n+1} over the unit ball $B_1(0) \subset \mathbb{R}^{n+1}$ in some detail.

2.1. Lemma. *Let π be the so-called Beltrami map*

$$(2.7) \quad \begin{aligned} \pi : \mathbb{H}^{n+1} &\rightarrow B_1(0) \subset \mathbb{R}^{n+1} \\ (x^0, x^i) &\rightarrow y = \left(\frac{x^i}{x^0}\right). \end{aligned}$$

Then π is a diffeomorphism such that, after introducing Euclidean polar coordinates (r, ξ) in $B_1(0)$, the hyperbolic metric can be expressed as

$$(2.8) \quad d\bar{s}^2 = \frac{r^2}{1-r^2} \left\{ \frac{1}{r^2(1-r^2)} dr^2 + \sigma_{ij} d\xi^i d\xi^j \right\}$$

or, if we define τ by

$$(2.9) \quad d\tau = \frac{1}{r\sqrt{1-r^2}} dr,$$

$$(2.10) \quad \begin{aligned} d\bar{s}^2 &= \frac{r^2}{1-r^2} \{d\tau^2 + \sigma_{ij} d\xi^i d\xi^j\} \\ &\equiv e^{2\psi} \{d\tau^2 + \sigma_{ij} d\xi^i d\xi^j\}; \end{aligned}$$

τ is uniquely determined up to an integration constant.

Proof. Writing the points in \mathbb{H}^{n+1} in the form (x^0, z) such that

$$(2.11) \quad -|x^0|^2 + |z|^2 = -1$$

we deduce

$$(2.12) \quad x^0 = \frac{1}{\sqrt{1-|y|^2}}, \quad y = \frac{z}{x^0},$$

hence

$$(2.13) \quad \pi^{-1}(y) = \left(\frac{1}{\sqrt{1-|y|^2}}, \frac{y}{\sqrt{1-|y|^2}} \right)$$

is a bijective mapping from $B_1(0)$ onto \mathbb{H}^{n+1} .

In polar coordinates $(y^\alpha) = (r, \xi^i)$, $1 \leq \alpha \leq n+1$,

$$(2.14) \quad x = \pi^{-1}(y) = \left(\frac{1}{\sqrt{1-r^2}}, \frac{r\xi}{\sqrt{1-r^2}} \right), \quad |\xi| = 1,$$

and the form (2.8) of the hyperbolic metric can be deduced from

$$(2.15) \quad \bar{g}_{\alpha\beta} = \langle x_\alpha, x_\beta \rangle, \quad 1 \leq \alpha, \beta \leq n+1. \quad \square$$

Now, let us denote the coordinates (τ, ξ^i) as usual by (x^α) , $0 \leq \alpha \leq n$, $\tau = x^0$, and let $(\bar{g}_{\alpha\beta})$ be the metric in (2.10).

Let $(\tilde{g}_{\alpha\beta})$ be the Euclidean metric in $B_1(0)$ in the coordinate system $(\tilde{x}^\alpha) = (\tilde{\tau}, x^i)$, such that

$$(2.16) \quad \begin{aligned} d\tilde{s}^2 &= \tilde{g}_{\alpha\beta} d\tilde{x}^\alpha d\tilde{x}^\beta = r^2 \{d\tilde{\tau}^2 + \sigma_{ij} d\xi^i d\xi^j\} \\ &\equiv e^{2\tilde{\psi}} \{d\tilde{\tau}^2 + \sigma_{ij} d\xi^i d\xi^j\}, \end{aligned}$$

where

$$(2.17) \quad d\tilde{\tau} = r^{-1} dr.$$

Writing $\tilde{\tau} = \varphi(\tau)$ we deduce

$$(2.18) \quad \frac{d\tilde{\tau}}{d\tau} = \frac{d\tilde{\tau}}{dr} \frac{dr}{d\tau} = \dot{\varphi} = \sqrt{1-r^2}.$$

Let $M \subset \mathbb{H}^{n+1}$ be an arbitrary closed, connected, strictly convex embedded hypersurface, then M is the boundary of a convex body \hat{M} . Without loss of generality we may assume that $x_0 = (1, 0) = \pi^{-1}(0)$ is an interior point of \hat{M} . Then M can be written as a graph in geodesic polar coordinates centered at x_0 , cf., e.g., [3, Section 4], or equivalently, as a graph over S^n in the coordinates (τ, x^i)

$$(2.19) \quad M = \text{graph } u = \{ \tau = u(x) : x \in S^n \}.$$

Because of (2.18) M can also be viewed as a graph \tilde{M} in $B_1(0)$ with respect to the Euclidean metric

$$(2.20) \quad \tilde{M} = \text{graph } \tilde{u} = \{ \tilde{\tau} = \tilde{u}(x) : x \in S^n \}.$$

Then we derive

$$(2.21) \quad \tilde{u} = \varphi(u).$$

Denote by (g_{ij}, h_{ij}) resp. $(\hat{g}_{ij}, \hat{h}_{ij})$ the metric and second fundamental form of M with respect to the ambient metrics $(\bar{g}_{\alpha\beta})$ resp. $(e^{-2\psi}\bar{g}_{\alpha\beta})$, and similarly let $(\tilde{g}_{ij}, \tilde{h}_{ij})$ resp. $(\hat{\tilde{g}}_{ij}, \hat{\tilde{h}}_{ij})$ be the geometric quantities of \tilde{M} with respect to the ambient metrics $(\tilde{g}_{\alpha\beta})$ resp. $(e^{-2\tilde{\psi}}\tilde{g}_{\alpha\beta})$.

Then we have

$$(2.22) \quad h_{ij}e^{-\psi} = \hat{h}_{ij} + \frac{d\psi}{d\tau}v^{-1}\hat{g}_{ij}$$

and

$$(2.23) \quad \tilde{h}_{ij}e^{-\tilde{\psi}} = \hat{\tilde{h}}_{ij} + \frac{d\tilde{\psi}}{d\tilde{\tau}}\tilde{v}^{-1}\hat{\tilde{g}}_{ij},$$

where

$$(2.24) \quad v^2 = 1 + \sigma^{ij}u_i u_j$$

$$(2.25) \quad \tilde{v}^2 = 1 + \sigma^{ij}\tilde{u}_i \tilde{u}_j,$$

see e.g., [7, Prop. 12.2.11].

Moreover, there holds

$$(2.26) \quad \hat{g}_{ij} = u_i u_j + \sigma_{ij}$$

and

$$(2.27) \quad \hat{h}_{ij} = -u_{ij}v^{-1},$$

where (u_{ij}) is the Hessian of u with respect to the metric (σ_{ij}) . Analogue formulas are valid for $\hat{\tilde{g}}_{ij}$ and $\hat{\tilde{h}}_{ij}$.

Hence we deduce

$$(2.28) \quad h_{ij}e^{-\psi} = -u_{ij}v^{-1} + \frac{d\psi}{d\tau}v^{-1}\hat{g}_{ij}$$

and

$$(2.29) \quad \tilde{h}_{ij}e^{-\tilde{\psi}} = -\tilde{u}_{ij}\tilde{v}^{-1} + \frac{d\tilde{\psi}}{d\tilde{\tau}}\tilde{v}^{-1}\hat{\tilde{g}}_{ij}.$$

Using the relations

$$(2.30) \quad \tilde{u}_i = \dot{\varphi}u_i,$$

and

$$(2.31) \quad \tilde{u}_{ij} = \dot{\varphi}u_{ij} + \ddot{\varphi}u_i u_j,$$

where

$$(2.32) \quad \dot{\varphi} = \sqrt{1 - r^2}, \quad \ddot{\varphi} = -r^2,$$

we obtain after some elementary calculations

$$(2.33) \quad \tilde{h}_{ij}\tilde{v} = (1 - r^2)h_{ij}v,$$

i.e., \tilde{M} is also strictly convex.

Moreover, let $u^i = \sigma^{ij}u_j$, then

$$(2.34) \quad \begin{aligned} \tilde{g}^{ij} &= r^{-2} \left\{ \sigma^{ij} - \frac{\dot{\varphi}^2}{\tilde{v}^2} u^i u^j \right\} \\ &\geq r^{-2} \left\{ \sigma^{ij} - \frac{1}{v^2} u^i u^j \right\}, \end{aligned}$$

since

$$(2.35) \quad \dot{\varphi}^2 = 1 - r^2 < 1$$

and we conclude

$$(2.36) \quad \tilde{h}_i^j \tilde{v} \geq h_i^j v,$$

hence,

$$(2.37) \quad \tilde{h}_i^j \geq h_i^j \frac{v}{\tilde{v}} \geq h_i^j.$$

Note also, that in points where $Du = 0$ there holds

$$(2.38) \quad \tilde{h}_i^j = h_i^j,$$

i.e., the principal curvatures are then identical.

Thus, we have proved

2.2. Lemma. *Let $M \subset \mathbb{H}^{n+1}$ be a closed, connected, strictly convex hypersurface, then the Beltrami map π maps M onto a closed strictly convex hypersurface $\tilde{M} \subset B_1(0)$. Moreover, expressing the normal vectors ν resp. $\tilde{\nu}$ of M resp. \tilde{M} in the common coordinate system (τ, ξ^i) yields that they are collinear.*

Proof. Only the last statement needs a verification. Up to a positive factor the covariant normal vector (ν_α) has the form

$$(2.39) \quad (\nu_\alpha) = (1, -u_i)$$

and $(\tilde{\nu}_\alpha)$, in the coordinate system (τ, ξ^i) ,

$$(2.40) \quad (\tilde{\nu}_\alpha) = \left(\frac{d\tilde{\tau}}{d\tau}, -\tilde{u}_i \right) = (\dot{\varphi}, -\dot{\varphi}u_i) = \dot{\varphi}(1, -u_i). \quad \square$$

2.3. Remark. The results of the preceding lemma can also be applied to a local embedding of a strictly convex hypersurface M that can be represented as a graph in geodesic polar coordinates centered in the Beltrami point $(1, 0)$ regardless which side of M the Beltrami point is facing.

3. HADAMARD'S THEOREM IN HYPERBOLIC SPACE

3.1. Theorem. *Let M_0 be a compact, connected n -dimensional manifold and*

$$(3.1) \quad x : M_0 \rightarrow \mathbb{H}^{n+1}$$

a strictly convex immersion of class C^2 , i.e., the second fundamental form with respect to any normal is always (locally) invertible, then the immersion is actually an embedding and $M = x(M_0)$ a strictly convex hypersurface that bounds a strictly convex body $\hat{M} \subset \mathbb{H}^{n+1}$. M and M_0 are moreover diffeomorphic to S^n and orientable.

Proof. Since we shall again employ the Beltrami map, we consider \mathbb{H}^{n+1} as a hypersurface in $\mathbb{R}^{n+1,1}$ and M as a codimension 2 immersed submanifold in $\mathbb{R}^{n+1,1}$, i.e.,

$$(3.2) \quad x : M_0 \rightarrow \mathbb{R}^{n+1,1}.$$

The Gaussian formula for M then looks like

$$(3.3) \quad x_{ij} = g_{ij}x - h_{ij}\tilde{x},$$

where g_{ij} is the induced metric, h_{ij} the second fundamental form of M considered as a hypersurface in \mathbb{H}^{n+1} , and \tilde{x} is the representation of the (exterior¹) normal vector $\nu = (\nu^\alpha)$ of M in $T(\mathbb{H}^{n+1})$ as a vector in $T(\mathbb{R}^{n+1,1})$.

Without loss of generality we may assume that the *Beltrami point* $(1, 0)$ does not belong to M , since the isometries of \mathbb{H}^{n+1} act transitively. Let π be the Beltrami map such that

$$(3.4) \quad \pi(1, 0) = 0 \in \mathbb{R}^{n+1}$$

and denote by φ its inverse

$$(3.5) \quad \varphi = \pi^{-1} : \mathbb{R}^{n+1} \rightarrow \mathbb{H}^{n+1} \subset \mathbb{R}^{n+1,1}.$$

Corresponding to the immersion $x = x(\xi)$ we then have an immersion $y = \pi \circ x$

$$(3.6) \quad y : M_0 \rightarrow \mathbb{R}^{n+1}.$$

Let $\tilde{M} \subset \mathbb{R}^{n+1}$ be its image and \tilde{g}_{ij} , \tilde{h}_{ij} , $\tilde{\nu} = (\tilde{\nu}^\alpha)$ its geometric quantities. We shall prove that \tilde{M} is an immersed, closed strictly convex hypersurface and hence an embedded hypersurface, due to Hadamard's theorem, cf. [16].

In view of the relation

$$(3.7) \quad x = \varphi \circ y$$

we shall then deduce that $x = x(\xi)$ is an embedding.

The inverse Beltrami map φ provides an embedding of \mathbb{H}^{n+1} in $\mathbb{R}^{n+1,1}$, i.e., we have the Gaussian formula

$$(3.8) \quad \varphi_{\alpha\beta} = \bar{g}_{\alpha\beta}\varphi,$$

¹Notice that for any closed, connected immersed hypersurface in \mathbb{H}^{n+1} an exterior normal vector can be unambiguously defined.

where $\bar{g}_{\alpha\beta}$ is the induced metric

$$(3.9) \quad \bar{g}_{\alpha\beta} = \langle \varphi_\alpha, \varphi_\beta \rangle.$$

Differentiating (3.7) covariantly with respect to the metric g_{ij} of M we obtain

$$(3.10) \quad \begin{aligned} x_{ij} &= \varphi_\alpha y_{ij}^\alpha + \varphi_{\alpha\beta} y_i^\alpha y_j^\beta \\ &= \varphi_\alpha y_{ij}^\alpha + \bar{g}_{\alpha\beta} y_i^\alpha y_j^\beta \varphi, \end{aligned}$$

in view of (3.8).

Indicate covariant derivatives with respect to the metric \tilde{g}_{ij} by a preceding semicolon such that

$$(3.11) \quad y_{;ij} = -\tilde{h}_{ij} \tilde{\nu},$$

then

$$(3.12) \quad y_{ij} = y_{;ij} - \{ \Gamma_{ij}^k - \tilde{\Gamma}_{ij}^k \} y_k,$$

hence, we derive from (3.10)

$$(3.13) \quad x_{ij} = \varphi_\alpha y_{;ij}^\alpha - \{ \Gamma_{ij}^k - \tilde{\Gamma}_{ij}^k \} y_k^\alpha \varphi_\alpha + \bar{g}_{\alpha\beta} y_i^\alpha y_j^\beta \varphi.$$

Let $\nu = (\nu^\alpha)$ be the exterior normal of M expressed in the coordinates (y^α) of \mathbb{R}^{n+1} , then the representation \tilde{x} of ν in $T(\mathbb{R}^{n+1,1})$ is given by

$$(3.14) \quad \tilde{x} = \varphi_\alpha \nu^\alpha$$

as can be easily checked.

From (3.3) and (3.13) we then deduce

$$(3.15) \quad h_{ij} = -\langle x_{ij}, \varphi_\gamma \nu^\gamma \rangle = \tilde{h}_{ij} \bar{g}_{\alpha\beta} \nu^\alpha \tilde{\nu}^\beta,$$

where we used

$$(3.16) \quad x_k = \varphi_\alpha y_k^\alpha$$

and

$$(3.17) \quad \langle x_k, \tilde{x} \rangle = 0.$$

Thus, it remains to prove that

$$(3.18) \quad \bar{g}_{\alpha\beta} \nu^\alpha \tilde{\nu}^\beta \neq 0 \quad \text{in } M_0.$$

So far we haven't used the fact $0 \notin \tilde{M}$, or equivalently, $(1, 0) \notin M$, but now we introduce polar coordinates (y^α) in \mathbb{R}^{n+1} such that $y^0 = r$ and distinguish two cases

$$(3.19) \quad \left\langle \frac{\partial}{\partial r}, \nu \right\rangle = 0$$

and

$$(3.20) \quad \left\langle \frac{\partial}{\partial r}, \nu \right\rangle \neq 0,$$

where the metric is the one in \mathbb{H}^{n+1} .

In Euclidean polar coordinates $(y^\alpha) = (r, \xi^i)$ the hyperbolic metric $(\bar{g}_{\alpha\beta})$ has been expressed in (2.8). Hence, if (3.19) is valid, we deduce

$$(3.21) \quad \nu^0 = 0,$$

and infer further, in view of (3.17),

$$(3.22) \quad \begin{aligned} 0 &= \langle x_i, \tilde{x} \rangle = \langle \varphi_\alpha, \varphi_\beta \rangle y_i^\alpha \nu^\beta \\ &= \bar{g}_{\alpha\beta} y_i^\alpha \nu^\beta = \frac{r^2}{1-r^2} \sigma_{kj} y_i^k \nu^j, \end{aligned}$$

from which we conclude that

$$(3.23) \quad \tilde{g}_{\alpha\beta} y_i^\alpha \nu^\beta = 0,$$

where $\tilde{g}_{\alpha\beta}$ is the Euclidean metric expressed in polar coordinates.

Thus, ν and $\tilde{\nu}$ are collinear, if (3.19) is valid.

On the other hand, if the assumption (3.20) is satisfied, then M , or more precisely, a local embedding of M_0 can be written as a graph in polar coordinates, i.e., we are in the situation where the results of Lemma 2.2 and the equations (2.39), (2.40) can be applied locally, cf. Remark 2.3, and we deduce again that $\nu, \tilde{\nu}$ are collinear.

Therefore, Hadamard's theorem yields that the immersion is actually an embedding and that M_0 , and hence M , is diffeomorphic to S^n . \square

4. THE GAUSS MAPS

Let $M \subset \mathbb{H}^{n+1}$ be a closed, connected, strictly convex hypersurface given by an embedding

$$(4.1) \quad x : M_0 \rightarrow M.$$

Considering M as a codimension 2 submanifold of $\mathbb{R}^{n+1,1}$ such that

$$(4.2) \quad x_{ij} = g_{ij}x - h_{ij}\tilde{x},$$

where $\tilde{x} \in T_x(\mathbb{R}^{n+1,1})$ represents the exterior normal vector $\nu \in T_x(\mathbb{H}^{n+1})$, we want to prove that the mapping

$$(4.3) \quad \tilde{x} : M_0 \rightarrow N$$

is an embedding of a strictly convex, closed, spacelike hypersurface \tilde{M} . We call this mapping the *Gauß map* of M .

First, we shall show that the Gauß map is injective. To prove this result we need the following lemma.

4.1. Lemma. *Let $M \subset \mathbb{H}^{n+1}$ be a closed, connected, strictly convex hypersurface and denote by \hat{M} its (closed) convex body. Let $x \in M$ be fixed and \tilde{x} be the corresponding outward normal vector, then*

$$(4.4) \quad \langle y, \tilde{x} \rangle \leq 0 \quad \forall y \in \hat{M}$$

and also strictly less than 0 unless $y = x$.

The preceding inequality also characterizes the points in \hat{M} , namely, let $y \in \mathbb{H}^{n+1}$ be such that

$$(4.5) \quad \langle y, \tilde{x} \rangle \leq 0 \quad \forall x \in M,$$

then $y \in \hat{M}$.

Proof. „(4.4)“ Let $y \in \text{int } \hat{M}$ be arbitrary and let $z = z(t)$, $0 \leq t \leq d$ be the unique geodesic in \mathbb{H}^{n+1} connecting y and x such that

$$(4.6) \quad z(0) = x \quad \wedge \quad z(d) = y$$

parametrized by arc length.

Viewing z as a curve in $\mathbb{R}^{n+1,1}$ the geodesic equation has the form

$$(4.7) \quad \ddot{z} \equiv \frac{D}{dt} \dot{z} = z.$$

If the coordinate system in $\mathbb{R}^{n+1,1}$ is Euclidean, the covariant derivatives are just ordinary derivatives.

It is well-known that the geodesic z is contained in \hat{M} and that

$$(4.8) \quad \langle \dot{z}(0), \tilde{x} \rangle < 0;$$

notice that, after introducing geodesic polar coordinates in \mathbb{H}^{n+1} centered in y , we have

$$(4.9) \quad \langle \dot{z}(0), \tilde{x} \rangle = -\left\langle \frac{\partial}{\partial r}, \nu \right\rangle$$

and hence is strictly negative, cf. [3, Section 4].

Thus, $\varphi(t) = \langle z(t), \tilde{x} \rangle$ satisfies the initial value problem

$$(4.10) \quad \ddot{\varphi} = \varphi, \quad \varphi(0) = 0, \quad \dot{\varphi}(0) < 0,$$

and is therefore equal to

$$(4.11) \quad \varphi(t) = -\lambda \sinh t, \quad \lambda > 0,$$

i.e.,

$$(4.12) \quad \varphi(t) < 0 \quad \forall t > 0.$$

Now, let $y \in M$, $y \neq x$, be arbitrary, and consider a sequence z_k of geodesics parametrized in the interval $0 \leq t \leq 1$, such that

$$(4.13) \quad z(0) = x \quad \wedge \quad z_k(1) \rightarrow y,$$

where $z_k(1) \in \text{int } \hat{M}$.

The geodesics z_k converge to a geodesic z connecting x and y . If

$$(4.14) \quad \langle \dot{z}(0), \tilde{x} \rangle < 0,$$

then the previous arguments are valid yielding

$$(4.15) \quad \langle y, \tilde{x} \rangle < 0.$$

On the other hand, the alternative

$$(4.16) \quad \langle y, \tilde{x} \rangle = 0$$

leads to a contradiction, since then the geodesic z would be part of the tangent space $T_x(M)$ which is impossible, cf. the considerations in [3] after the equation (4.17).

„ $y \in \hat{M}$ “ Suppose now that $y \in \mathbb{H}^{n+1}$ satisfies (4.5), and assume by contradiction that $y \in \mathbb{C}\hat{M}$. Pick an arbitrary $x_0 \in \text{int}\hat{M}$ and let $z = z(t)$, $0 \leq t \leq d$, be the geodesic joining x_0 and y parameterized by arc length, such that $z(0) = x_0$ and $z(d) = y$. The geodesic intersects M in a unique point x , $x = z(t_1)$, $0 < t_1 < d$.

Define

$$(4.17) \quad \varphi(t) = \langle z(t), \tilde{x} \rangle$$

and let $0 \leq t_0 \leq d$ be such that

$$(4.18) \quad \varphi(t_0) = \sup\{\varphi(t) : 0 \leq t \leq d\}.$$

We now distinguish two cases. First, we assume $\varphi(t_0) > 0$, then there must hold $0 < t_0 < d$ and $\dot{\varphi}(t_0) = 0$. Thus φ satisfies the initial value problem

$$(4.19) \quad \ddot{\varphi} = \varphi, \quad \varphi(t_0) > 0, \quad \dot{\varphi}(t_0) = 0,$$

and must therefore be equal to

$$(4.20) \quad \varphi(t) = \lambda \cosh(t - t_0), \quad \lambda > 0,$$

which is a contradiction, since $\varphi(0) < 0$.

Hence, we must have $\varphi(t_0) = 0$ and we may choose $t_0 = t_1$, i.e., there holds $\dot{\varphi}(t_1) = 0$, which is a contradiction too, because of the inequality (4.8), which now reads $\dot{\varphi}(t_1) > 0$.

Therefore we have proved $y \in \hat{M}$. □

4.2. Theorem. *Let $x : M_0 \rightarrow M \subset \mathbb{H}^{n+1}$ be the embedding of a closed, connected, strictly convex hypersurface, then the Gauß map defined in (4.3) is injective, where we identify $\mathbb{R}^{n+1,1}$ with its individual tangent spaces.*

Proof. We again assume M to be a codimension 2 submanifold in $\mathbb{R}^{n+1,1}$. Suppose there would be two points $p_1 \neq p_2$ in M_0 such that

$$(4.21) \quad \tilde{x}(p_1) = \tilde{x}(p_2),$$

then the function

$$(4.22) \quad \varphi(y) = \langle y, \tilde{x}(p_1) \rangle$$

would vanish in the points $x(p_1)$ as well as $x(p_2)$ contrary to the results of Lemma 4.1. □

4.3. Lemma. *As a submanifold of codimension 2 M satisfies the Weingarten equations*

$$(4.23) \quad \tilde{x}_i = h_i^k x_k$$

for the normal \tilde{x} and also

$$(4.24) \quad x_i = g_i^k x_k$$

for the normal x .

Proof. We only have to prove the non-trivial Weingarten equation.

First we infer from

$$(4.25) \quad \langle x, \tilde{x} \rangle = 0$$

that

$$(4.26) \quad 0 = \langle x_i, \tilde{x} \rangle + \langle x, \tilde{x}_i \rangle = \langle x, \tilde{x}_i \rangle.$$

Furthermore, there holds

$$(4.27) \quad 0 = \langle \tilde{x}, \tilde{x}_i \rangle,$$

since $\langle \tilde{x}, \tilde{x} \rangle = 1$. Hence, we deduce

$$(4.28) \quad \tilde{x}_i = a_i^k x_k.$$

Differentiating the relation $\langle x_j, \tilde{x} \rangle = 0$ covariantly we obtain

$$(4.29) \quad \langle \tilde{x}_j, x_i \rangle = h_{ij}$$

and we infer (4.23) in view of (4.28). \square

We can now prove

4.4. Theorem. *Let $x : M_0 \rightarrow M \subset \mathbb{H}^{n+1}$ be a closed, connected, strictly convex hypersurface of class C^m , $m \geq 3$, then the Gauß map \tilde{x} in (4.3) is the embedding of a closed, spacelike, achronal, strictly convex hypersurface $\tilde{M} \subset N$ of class C^{m-1} .*

Viewing \tilde{M} as a codimension 2 submanifold in $\mathbb{R}^{n+1,1}$, its Gaussian formula is

$$(4.30) \quad \tilde{x}_{ij} = -\tilde{g}_{ij}\tilde{x} + \tilde{h}_{ij}x,$$

where \tilde{g}_{ij} , \tilde{h}_{ij} are the metric and second fundamental form of the hypersurface $\tilde{M} \subset N$, and $x = x(\xi)$ is the embedding of M which also represents the future directed normal vector of \tilde{M} . The second fundamental form \tilde{h}_{ij} is defined with respect to the future directed normal vector, where the time orientation of N is inherited from $\mathbb{R}^{n+1,1}$.

The second fundamental forms of M , \tilde{M} and the corresponding principal curvatures κ_i , $\tilde{\kappa}_i$ satisfy

$$(4.31) \quad h_{ij} = \tilde{h}_{ij} = \langle \tilde{x}_i, x_j \rangle$$

and

$$(4.32) \quad \tilde{\kappa}_i = \kappa_i^{-1}.$$

Proof. (i) From the Weingarten equation (4.23) we infer

$$(4.33) \quad \tilde{g}_{ij} = \langle \tilde{x}_i, \tilde{x}_j \rangle = h_i^k h_{kj}$$

is positive definite, hence $\tilde{x} = \tilde{x}(\xi)$ is an embedding of a closed, connected spacelike hypersurface, where we also used Theorem 4.2.

Since N is simply connected, we conclude further that \tilde{M} is achronal and thus can be written as a graph over the Cauchy hypersurface $\{0\} \times S^n$ which we identify with S^n

$$(4.34) \quad \tilde{M} = \text{graph } \tilde{u}|_{S^n} \subset N,$$

cf. [14, p. 427] and [6, Prop. 2.5].

(ii) The pair (x, \tilde{x}) satisfies

$$(4.35) \quad \langle x, \tilde{x} \rangle = 0$$

and we claim that x is the future directed normal vector of \tilde{M} in \tilde{x} , where as usual we identify the normal vector $\tilde{\nu} = (\tilde{\nu}^\alpha) \in T_{\tilde{x}}(N)$ with its embedding in $T_{\tilde{x}}(\mathbb{R}^{n+1,1})$.

Differentiating (4.35) covariantly and using the fact that \tilde{x} is a normal vector for M we deduce

$$(4.36) \quad 0 = \langle x, \tilde{x}_i \rangle,$$

i.e., \tilde{x} and x span the normal space of the codimension 2 submanifold \tilde{M} . By the very definition of \mathbb{H}^{n+1} x is a future directed vector in $\mathbb{R}^{n+1,1}$.

Let us define the second fundamental form \tilde{h}_{ij} of $\tilde{M} \subset N$ with respect to the future directed normal vector $\tilde{\nu} \in T_{\tilde{x}}(N)$, then the codimension 2 Gaussian formula is exactly (4.30) because of (4.36).

Differentiating the Weingarten equation (4.23) covariantly with respect to the metric \tilde{g}_{ij} and indicating the covariant derivatives with respect to \tilde{g}_{ij} by a semi-colon and those with respect to g_{ij} simply by indices, we obtain

$$(4.37) \quad \tilde{x}_{;ij} = h_{i;j}^k x_k + h_i^k x_{;kj}$$

and we deduce further

$$(4.38) \quad \tilde{h}_{ij} = -\langle \tilde{x}_{;ij}, x \rangle = -h_i^k \langle x_{kj}, x \rangle = h_i^k g_{kj} = h_{ij}.$$

On the other hand, we infer from (4.36)

$$(4.39) \quad \tilde{h}_{ij} = -\langle \tilde{x}_{;ij}, x \rangle = \langle \tilde{x}_i, x_j \rangle$$

which proves (4.31).

The last relation (4.32) follows from (4.38) and (4.33). \square

We can also define a Gauß map from strictly convex, connected, spacelike hypersurfaces $\tilde{M} \subset N$ into \mathbb{H}^{n+1} such that the two Gauß maps are inverse to each other.

4.5. Theorem. *Let $\tilde{M} \subset N$ be a closed, connected, spacelike, strictly convex, embedded hypersurface of class C^m , $m \geq 3$, such that, when viewed as a codimension 2 submanifold in $\mathbb{R}^{n+1,1}$, its Gaussian formula is*

$$(4.40) \quad \tilde{x}_{ij} = -\tilde{g}_{ij} \tilde{x} + \tilde{h}_{ij} x,$$

where $\tilde{x} = \tilde{x}(\xi)$ is the embedding, x the future directed normal vector, and \tilde{g}_{ij} , \tilde{h}_{ij} the induced metric and the second fundamental form of the hypersurface in N . Then we define the Gauß map as $x = x(\xi)$

$$(4.41) \quad x : \tilde{M} \rightarrow \mathbb{H}^{n+1} \subset \mathbb{R}^{n+1,1}.$$

The Gauß map is the embedding of a closed, connected, strictly convex hypersurface M in \mathbb{H}^{n+1} .

Let g_{ij} , h_{ij} be the induced metric and second fundamental form of M , then, when viewed as a codimension 2 submanifold, M satisfies the relations

$$(4.42) \quad x_{ij} = g_{ij}x - h_{ij}\tilde{x},$$

$$(4.43) \quad h_{ij} = \tilde{h}_{ij} = \langle x_i, \tilde{x}_j \rangle,$$

and

$$(4.44) \quad \kappa_i = \tilde{\kappa}_i^{-1},$$

where κ_i , $\tilde{\kappa}_i$ are the corresponding principal curvatures.

Proof. The fact that $x = x(\xi)$ is the immersion of a closed, connected, strictly convex hypersurface M satisfying the relations (4.42), (4.43), and (4.44) follows along the lines of the proof of the previous theorem.

Using Theorem 3.1 we then deduce that the immersion is an embedding. \square

Combining the two theorems, looking especially at the Gaussian formulas (4.30) and (4.42), we immediately conclude that the Gauß maps are inverse to each other, i.e., if we start with a closed, strictly convex hypersurface $M \subset \mathbb{H}^{n+1}$, apply the Gauß map to obtain a spacelike, strictly convex hypersurface $\tilde{M} \subset N$, and then apply the second Gauß map, then we return to M with a pointwise equality.

Denoting the two Gauß maps simply by a tilde, this can be expressed in the form

$$(4.45) \quad x = \tilde{\tilde{x}},$$

or, equivalently, in the form of a commutative diagram

$$(4.46) \quad \begin{array}{ccc} M & \xrightarrow{\tilde{}} & \tilde{M} \\ & \searrow \text{id} & \swarrow \tilde{} \\ & & M \end{array}$$

Before we give an equivalent characterization of the images of the Gauß maps, let us show that the images of strictly convex hypersurfaces by the Gauß maps are as smooth as the original hypersurfaces.

4.6. Theorem. *Let $M \subset \mathbb{H}^{n+1}$ be a closed, connected, strictly convex hypersurface of class $C^{m,\alpha}$, $m \geq 2$, $0 \leq \alpha \leq 1$, then $\tilde{M} \subset N$, its image under the Gauß map is also of class $C^{m,\alpha}$.*

The corresponding regularity result is also valid, if we start with a closed, spacelike, connected, strictly convex hypersurface in N and use the Gauß map to embed it into \mathbb{H}^{n+1} .

Proof. We only consider the case when we apply the Gauß map to $M \subset \mathbb{H}^{n+1}$. Moreover, without loss of generality we shall also assume that the Beltrami point $(1, 0)$ is not part of M .

(i) First, let us assume that $m \geq 3$ and $0 \leq \alpha \leq 1$. The Gauß map is then of class $C^{m-1,\alpha}$, i.e., \tilde{M} is of class $C^{m-1,\alpha}$. Here, we use the coordinates (ξ^i) for M also as coordinates for \tilde{M} . The metric \tilde{g}_{ij} and the Christoffel symbols of \tilde{M} are then of class $C^{m-2,\alpha}$ resp. $C^{m-3,\alpha}$, while the second fundamental form \tilde{h}_{ij} is of class $C^{m-2,\alpha}$, in view of (4.31).

Representing now \tilde{M} as a graph over S^n ,

$$(4.47) \quad \tilde{M} = \text{graph } u|_{S^n}$$

in conformal coordinates, i.e., we use the coordinates defined in the formulas (2.3) to (2.6) denoting them this time, however, by (τ, x^i) instead of (τ, ξ^i) , since (ξ^i) are supposed to be given coordinates for M .

Notice that the transformation $(x^i(\xi^k))$ is a diffeomorphism of class $C^{m-1,\alpha}$, since the underlying polar coordinates (x^0, r, x^i) , defined in (2.3), also cover that part of $\mathbb{R}^{n+1,1}$ that contains M , due to our assumption at the beginning of the proof. Hence, expressing the Gauß map in this ambient coordinate system

$$(4.48) \quad \tilde{x}(\xi) = (x^0(\xi), r(\xi), x^i(\xi)),$$

where

$$(4.49) \quad r = \sqrt{1 + |x^0|^2},$$

we deduce

$$(4.50) \quad \begin{aligned} \tilde{g}_{ij} &= \langle \tilde{x}_i, \tilde{x}_j \rangle = -x_i^0 x_j^0 + r_i r_j + r^2 \sigma_{kl} x_i^k x_j^l \\ &= -\frac{1}{1 + |x^0|^2} x_i^0 x_j^0 + (1 + |x^0|^2) \sigma_{kl} x_i^k x_j^l \end{aligned}$$

in view of (2.3), proving that the Jacobian (x_i^k) is invertible.

Thus, we conclude that the second fundamental form \tilde{h}_{ij} expressed in the new coordinates (x^i) is still of class $C^{m-2,\alpha}$.

We want to express the covariant derivatives u_{ij} of u with respect to the metric σ_{ij} in terms of \tilde{h}_{ij} to deduce that u_{ij} is of class $C^{m-1,\alpha}$, and hence $u \in C^{m,\alpha}(S^n)$.

To achieve this we define a new metric $\hat{g}_{\alpha\beta}$ in the ambient space

$$(4.51) \quad \hat{g}_{\alpha\beta} = e^{-2\psi} \tilde{g}_{\alpha\beta},$$

where $\tilde{g}_{\alpha\beta}$ is the metric in (2.5). Let \hat{g}_{ij} , \hat{h}_{ij} and $\hat{\nu}$ be the obvious geometric quantities of \tilde{M} with respect to the new metric, then there holds

$$(4.52) \quad h_{ij}e^{-\psi} = \hat{h}_{ij} + \psi_\alpha \hat{\nu}^\alpha \hat{g}_{ij}$$

cf. (2.23), where we already used this formula.

On the other hand, \hat{h}_{ij} can be expressed in terms of the Hessian $u_{;ij}$ of u with respect to the metric σ_{ij} , namely,

$$(4.53) \quad \hat{h}_{ij} = u_{;ij}v^{-1},$$

i.e.,

$$(4.54) \quad h_{ij}e^{-\psi} = u_{;ij} + \psi_\alpha \hat{\nu}^\alpha (-u_i u_j + \sigma_{ij}),$$

hence, $u_{;ij}$ is of class $C^{m-2,\alpha}$.

(ii) The case $m = 2$ and $0 \leq \alpha \leq 1$ follows by approximation and the uniform $C^{2,\alpha}$ -estimates. Notice that the approximating second fundamental forms will converge in C^0 . \square

4.7. Definition. (i) Let $M \subset \mathbb{H}^{n+1}$ be a closed, connected, strictly convex hypersurface, then we define its *polar set* $M^* \subset N$ by

$$(4.55) \quad M^* = \{y \in N : \sup_{x \in M} \langle x, y \rangle = 0\},$$

where the scalar product is the scalar product in $\mathbb{R}^{n+1,1}$ and x, y are Euclidean coordinates.

(ii) A similar definition holds, if $M \subset N$ is a spacelike, closed, connected, strictly convex hypersurface $M \subset N$, then

$$(4.56) \quad M^* = \{y \in \mathbb{H}^{n+1} : \sup_{x \in M} \langle x, y \rangle = 0\}.$$

4.8. Theorem. *The polar sets agree with the images of the Gauß maps.*

Proof. Again we only consider the case $M \subset \mathbb{H}^{n+1}$.

In view of Lemma 4.1 there holds

$$(4.57) \quad \tilde{M} \subset M^*.$$

On the other hand, let $y \in M^*$ and $x \in M$ be such that

$$(4.58) \quad \langle x, y \rangle = 0.$$

Then we deduce, after introducing local coordinates in M ,

$$(4.59) \quad \langle x_i, y \rangle = 0$$

and

$$(4.60) \quad \langle x_{ij}, y \rangle \leq 0,$$

where the derivatives are covariant derivatives with respect to the induced metric g_{ij} of M being viewed as a codimension 2 submanifold.

Combining (4.58) and (4.59) we infer

$$(4.61) \quad y = \pm \tilde{x},$$

but because of (4.42) and (4.60) we deduce $y = \tilde{x}$. \square

Let us conclude the section with

4.9. Theorem. *The Gauß maps provide a bijective relation between the connected, closed, strictly convex hypersurfaces $M \subset \mathbb{H}^{n+1}$ having the Beltrami point in the interior of their convex bodies and the spacelike, closed, connected, strictly convex hypersurfaces $\tilde{M} \subset N_+$, where*

$$(4.62) \quad N_+ = \{x \in N : x^0 > 0\}.$$

The geodesic spheres with center in the Beltrami point are mapped onto the coordinate slices $\{x^0 = \text{const}\}$.

Proof. (i) Let $M \subset \mathbb{H}^{n+1}$ be closed, strictly convex such that $p_0 \in \text{int}\hat{M}$, where $p_0 = (1, 0, \dots, 0) \in \mathbb{R}^{n+1,1}$. According to Lemma 2.1, $\mathbb{H}^{n+1} \subset \mathbb{R}^{n+1,1}$ can be written as the embedding of \mathbb{R}^{n+1} via the inverse $\varphi = \pi^{-1}$ of the Beltrami map π , and M can be represented as $M = \text{graph } u|_{S^n}$ in geodesic polar coordinates centered in p_0 , or more precisely, in the coordinates (τ, ξ^i) .

A moment's reflection reveals that the Gauß map of M is given by

$$(4.63) \quad \tilde{x} = \varphi_\alpha \nu^\alpha, \quad \nu \in T_x(\mathbb{H}^{n+1}),$$

where ν is the exterior normal. In geodesic polar coordinates the normal ν is given by

$$(4.64) \quad (\nu^\alpha) = \tilde{v} e^{-\psi} (1, -u^i),$$

where $\tilde{v} = v^{-1}$ and $u^i = \sigma^{ij} u_j$; notice that the metric in \mathbb{H}^{n+1} is expressed as in (2.10).

Hence, we deduce from (2.14)

$$(4.65) \quad \tilde{x}^0 = \frac{r^2}{(1-r^2)^{3/2}} \tilde{v} e^{-\psi} = \frac{r}{\sqrt{1-r^2}} \tilde{v} > 0,$$

i.e., $\tilde{M} \subset N_+$.

If M is a geodesic sphere, then we deduce from (4.63) and (4.64) that it is mapped onto a coordinate slice in N_+ .

(ii) To prove the inverse relation, consider a spacelike, closed, connected, strictly convex hypersurface $M \subset N_+$. Assuming the coordinate system in (2.4), (2.5), N_+ can be viewed as the embedding of $\mathbb{R}^{n+1} \setminus \bar{B}_1(0)$ in $\mathbb{R}^{n+1,1}$ via the map

$$(4.66) \quad x = \varphi(r, \xi) = (\sqrt{r^2 - 1}, r\xi), \quad \xi \in S^n.$$

The Gauß map from M into \mathbb{H}^{n+1} can then be expressed as

$$(4.67) \quad \tilde{x} = \varphi_\alpha \nu^\alpha,$$

where

$$(4.68) \quad \nu = \tilde{\nu} e^{-\psi}(1, u^i)$$

is the future directed normal vector in $x \in M$. Let $\tilde{M} \subset \mathbb{H}^{n+1}$ be its image. Then we have to show that the Beltrami point p_0 is an interior point of the corresponding convex body, or equivalently, that

$$(4.69) \quad \langle p_0, x \rangle < 0 \quad \forall x \in M,$$

in view of the second part of Lemma 4.1. But we immediately deduce

$$(4.70) \quad \langle p_0, x \rangle = -\sqrt{r^2 - 1},$$

in view of (4.66).

Again we conclude from (4.67) that coordinate slices are mapped onto geodesic spheres. \square

5. CURVATURE FLOW

Let us now consider the problem of finding a solution of

$$(5.1) \quad F|_M = f(\nu),$$

where F is a curvature function defined in the open positive cone $\Gamma_+ \subset \mathbb{R}^n$, $0 < f$ is a function defined in the normal space of M , and $M \subset \mathbb{H}^{n+1}$ is a closed, connected, strictly convex hypersurface yet to be determined.

Using the results of the previous section, especially Theorem 4.4 and Theorem 4.5, we can reformulate the problem equivalently by assuming that $0 < f \in C^{2,\alpha}(N)$, $0 < \alpha < 1$, is given and a closed, strictly convex hypersurface $M \subset \mathbb{H}^{n+1}$ is to be found satisfying

$$(5.2) \quad F|_M = f(\tilde{x}) \quad \forall x \in M,$$

where $x \rightarrow \tilde{x}$ is the Gauß map corresponding to M .

Let $\tilde{M} \subset N$ be the image of M under the Gauß map, which is identical with the polar M^* of M , and let \tilde{F} be the inverse of F , i.e.,

$$(5.3) \quad \tilde{F}(\kappa_i) = \frac{1}{F(\kappa_i^{-1})},$$

then the equation (5.2) is equivalent to

$$(5.4) \quad \tilde{F}|_{\tilde{M}} = f^{-1}(\tilde{x}) \quad \forall \tilde{x} \in \tilde{M},$$

in view of (4.32), where now the right-hand side depends on the points $\tilde{x} \in \tilde{M}$, and $\tilde{M} \subset N$ is a closed, spacelike, connected strictly convex hypersurface in the de Sitter space N with curvature $K_N = 1$.

We solved problems of this kind in [6, Theorem 0.2] assuming barrier conditions and some additional hypotheses. In that paper we denoted the curvature function, the right-hand side and the hypersurface by F , f , and M , which would correspond to the present notation \tilde{F} , f^{-1} , \tilde{M} .

Let us stick to the last notation just long enough to formulate the condition for \tilde{F} , namely, we assume that \tilde{F} is of class (K^*) , where the class (K^*) is defined in [6, Definition 1.6] as

5.1. Definition. A symmetric curvature function $F \in C^{2,\alpha}(\Gamma_+) \cap C^0(\bar{\Gamma}_+)$ of class $(K)^2$ is said to be of class (K^*) , if there exists $0 < \epsilon_0 = \epsilon_0(F)$ such that

$$(5.5) \quad \epsilon_0 F H \leq F^{ij} h_{ik} h_j^k$$

for any positive symmetric tensor (h_{ij}) . Here (h_{ij}) and a Riemannian metric (g_{ij}) should be defined in a given tensor space $T^{0,2}$, and H stands for the trace of (h_{ij})

$$(5.6) \quad H = g^{ij} h_{ij}.$$

5.2. Remark. Functions F that can be written as

$$(5.7) \quad F = GK^a, \quad a > 0,$$

where K is the Gaussian curvature and G an arbitrary function of class (K) , including the case $G = 1$, which doesn't belong to (K) , are of class (K^*) , cf. [6, Proposition 1.9].

Thus, we shall solve the original problem (5.2) for curvature functions F satisfying the requirement that their inverses $\tilde{F} \in (K^*)$.

Notice that in case $F = K$ there holds

$$(5.8) \quad F = \tilde{F} \in (K^*).$$

We shall also assume without loss of generality that F is homogeneous of degree 1, and hence concave, cf. [8, Lemma 1.2].

Now that we have formulated the condition for \tilde{F} , let us switch notations to enhance the readability of the text and to simplify the comparison with former results, and let us rewrite the equation (5.4) in the form

$$(5.9) \quad F|_M = f(x) \quad \forall x \in M,$$

where $F \in (K^*)$, $0 < f$ is defined in N and $M \subset N$ is a closed, spacelike, connected, strictly convex hypersurface, where its second fundamental form is defined with respect to the future directed normal, in contrast to our default convention to consider the past directed normal.

In order to make the comparison with former results and techniques easier, we therefore switch the light cone, so that the future directed normal is now past directed, and replace the time function τ in (2.6) by $-\tau$ without changing the notation, i.e., x^0 is still the time function inherited from $\mathbb{R}^{n+1,1}$, but

$$(5.10) \quad \frac{dx^0}{d\tau} = -(1 + |x^0|^2),$$

²For a definition of the class (K) see [6, Definition 1.1].

and the coordinate slices with positive curvature are now contained in $\{\tau < 0\}$.

We want to solve equation (5.9). For technical reasons, it is convenient to solve instead the equivalent equation

$$(5.11) \quad \Phi(F)|_M = \Phi(f),$$

where Φ is a real function defined on \mathbb{R}_+ such that

$$(5.12) \quad \dot{\Phi} > 0 \quad \text{and} \quad \ddot{\Phi} \leq 0.$$

For notational reasons, let us abbreviate

$$(5.13) \quad \tilde{f} = \Phi(f).$$

We also point out that we may—and shall—assume without loss of generality that F is homogeneous of degree 1.

To solve (5.11) we look at the evolution problem

$$(5.14) \quad \begin{aligned} \dot{x} &= (\Phi - \tilde{f})\nu, \\ x(0) &= x_0, \end{aligned}$$

where x_0 is an embedding of an initial strictly convex, compact, space-like hypersurface M_0 , $\Phi = \Phi(F)$, and F is evaluated at the principal curvatures of the flow hypersurfaces $M(t)$, or, equivalently, we may assume that F depends on the second fundamental form (h_{ij}) and the metric (g_{ij}) of $M(t)$; $x(t)$ is the embedding of $M(t)$, and ν is the past directed normal of the flow hypersurfaces $M(t)$.

This is a parabolic problem, so short-time existence is guaranteed—the proof in the Lorentzian case is identical to that in the Riemannian case, cf. [3, p. 622]—, and under suitable assumptions, which we are going to formulate in a moment, we shall be able to prove that the solution exists for all time and converges to a stationary solution if t goes to infinity.

In N we consider an open, connected, precompact set Ω that is bounded by two *achronal*, connected, spacelike hypersurfaces M_1 and M_2 , where M_1 is supposed to lie in the *past* of M_2 .

We assume that $0 < f \in C^{2,\alpha}(\bar{\Omega})$, $0 < \alpha < 1$, and that the boundary components M_i act as barriers for (F, f) .

5.3. Definition. M_2 is an *upper barrier* for (F, f) , if M_2 is strictly convex and satisfies

$$(5.15) \quad F|_{M_2} \geq f,$$

and M_1 is a *lower barrier* for (F, f) , if at the points $\Sigma \subset M_1$, where M_1 is strictly convex, there holds

$$(5.16) \quad F|_{\Sigma} \leq f.$$

Σ may be empty.

To simplify some calculations that are to follow, we introduce an eigen time coordinate system in N , i.e., we write the metric in the form

$$(5.17) \quad d\bar{s}^2 = -d\tau^2 + \cosh^2 \tau \sigma_{ij} d\xi^i d\xi^j,$$

where σ_{ij} is the standard metric of S^n . The time function τ is globally defined, and due to our convention the uniform convex slices are contained in $\{\tau < 0\}$.

This preceding relation can be immediately deduced from (2.5) and (2.6). The special form of the metric with $\cosh^2 \tau$ is of no importance. The crucial facts are that N has constant curvature, $\frac{\partial}{\partial \tau}$ is a timelike unit vector field, and the coordinate slices $\{\tau = \text{const}\}$ are totally umbilic.

Notice also that, if $M = \text{graph } u$ is a spacelike hypersurface, the previously defined quantities v and \tilde{v} are identical to those defined in the new coordinate system

$$(5.18) \quad v^2 = 1 - \bar{g}^{ij} u_i u_j, \quad \bar{g}_{ij} = \cosh^2 \tau \sigma_{ij}.$$

However, when applying the formulas in Section 1 one should observe that in the present coordinate system the terms in equation (1.8) should read

$$(5.19) \quad \psi = 0 \quad \wedge \quad \sigma_{ij} = \bar{g}_{ij}.$$

We now consider the evolution problem (5.14) with $M_0 = M_2$. Then the flow exists in a maximal time interval $I = [0, T^*)$, $0 < T^* \leq \infty$, and, as we have proved in [6, Section 4], there holds

5.4. Lemma. *During the evolution the flow hypersurfaces stay inside $\bar{\Omega}$. The hypersurfaces $M(t)$ can be written as graphs over S^n*

$$(5.20) \quad M(t) = \text{graph } u(t, \cdot),$$

such that, if the barriers are expressed as $M_i = \text{graph } u_i$, $i = 1, 2$, we have

$$(5.21) \quad u_1 \leq u \leq u_2$$

and the quantity

$$(5.22) \quad \tilde{v} = \frac{1}{\sqrt{1 - |Du|^2}}$$

is uniformly bounded for all $t \in I$.

Moreover, the initial inequality $F \geq f$ is valid throughout the evolution, which can be equivalently formulated as

$$(5.23) \quad \Phi \geq \tilde{f}.$$

Let us now look at the evolution equations satisfied by u , \tilde{v} , and h_i^j .

5.5. Lemma. *Let $M(t) = \text{graph } u(t, \cdot)|_{S^n}$ be the flow hypersurfaces, then u satisfies the parabolic equation*

$$(5.24) \quad \dot{u} - \dot{\Phi} F^{ij} u_{ij} = -\tilde{v}(\Phi - \tilde{f}) + \tilde{v} \dot{\Phi} F - \dot{\Phi} F^{ij} \bar{h}_{ij},$$

where

$$(5.25) \quad \dot{\Phi} = \frac{d\Phi(r)}{dr} \quad \wedge \quad \dot{u} = \frac{du}{dt}.$$

Proof. These equations immediately follow from the relations (1.17), (1.18) and (5.14) by observing that

$$(5.26) \quad \dot{u} = \dot{x}^0.$$

Notice that \dot{u} is the total time derivative, where „time“ is just the usual name for the flow parameter. \square

5.6. Lemma. *The quantity \tilde{v} satisfies the evolution equation*

$$(5.27) \quad \begin{aligned} \dot{\tilde{v}} - \dot{\Phi} F^{ij} \tilde{v}_{ij} &= -\dot{\Phi} F^{ij} h_{ik} h_j^k \tilde{v} - [(\Phi - \tilde{f}) - \dot{\Phi} F] \bar{h}_{ij} \tilde{u}^i \tilde{u}^j \tilde{v}^2 \\ &+ 2\dot{\Phi} F^{ij} h_i^k \bar{h}_{ki} - \bar{\kappa}^2 \dot{\Phi} F^{ij} \bar{g}_{ij} \tilde{v} - 2\bar{\kappa}^2 \dot{\Phi} F^{ij} u_i u_j \tilde{v} \\ &+ K_N \dot{\Phi} F^{ij} u_i u_j \tilde{v} + f_{\beta} x_i^{\beta} u^i, \end{aligned}$$

where $\bar{\kappa}$ is the principal curvature of the slices $\{\tau = \text{const}\}$,

$$(5.28) \quad \tilde{u}^i = \bar{g}^{ij} u_j$$

and $K_N = 1$ for the de Sitter space-time.

Proof. Let $(\eta_{\alpha}) = (-1, 0, \dots, 0)$ be the covariant vector field representing $-\frac{\partial}{\partial \tau}$. Differentiating $\tilde{v} = \eta_{\alpha} \nu^{\alpha}$ covariantly with respect to the induced metric of M , where (ν^{α}) is the past directed normal, we obtain

$$(5.29) \quad \begin{aligned} \dot{\tilde{v}} - \dot{\Phi} F^{ij} \tilde{v}_{ij} &= -\dot{\Phi} F^{ij} h_{ik} h_j^k \tilde{v} + [(\Phi - \tilde{f}) - \dot{\Phi} F] \eta_{\alpha\beta} \nu^{\alpha} \nu^{\beta} \\ &- 2\dot{\Phi} F^{ij} h_j^k x_i^{\alpha} x_k^{\beta} \eta_{\alpha\beta} - \dot{\Phi} F^{ij} \eta_{\alpha\beta\gamma} x_i^{\beta} x_j^{\gamma} \nu^{\alpha} \\ &- \dot{\Phi} F^{ij} \bar{R}_{\alpha\beta\gamma\delta} \nu^{\alpha} x_i^{\beta} x_k^{\gamma} \eta_{\epsilon} x_l^{\delta} \eta_{\epsilon}^{\epsilon} g^{kl} \\ &- \tilde{f}_{\beta} x_i^{\beta} x_k^{\alpha} \eta_{\alpha} g^{ik}, \end{aligned}$$

where the ambient space can be a general Lorentzian manifold, cf. [6, Lemma 4.4]; however, in the general case the definition of η has to be adjusted, since it has to be a unit vector field.

Now, if the ambient space is a space of constant curvature K_N , the term containing the Riemannian curvature tensor vanishes, and we shall show that the crucial term

$$(5.30) \quad -F^{ij} \eta_{\alpha\beta\gamma} x_i^{\beta} x_j^{\gamma} \nu^{\alpha}$$

can be expressed as claimed.

First we observe that the second fundamental form \bar{h}_{ij} of the coordinate slices $\{\tau = \text{const}\}$ is given by

$$(5.31) \quad \bar{h}_{ij} = -\frac{1}{2} \dot{g}_{ij}.$$

Secondly, the vector field (η_{α}) is a gradient field, namely,

$$(5.32) \quad (\eta_{\alpha}) = \text{grad } \varphi$$

with

$$(5.33) \quad \varphi(\tau, x^i) = -\tau.$$

Since $D\varphi$ is a unit vector field, we have

$$(5.34) \quad \varphi_{\alpha\beta}\varphi^\alpha = \varphi_{\beta\alpha}\varphi^\alpha = 0.$$

The restriction of φ to a coordinate slice is constant, hence, differentiating φ covariantly with respect to the induced metric \bar{g}_{ij} , we deduce

$$(5.35) \quad \begin{aligned} 0 = \varphi_{ij} &= \varphi_\alpha \bar{x}_{ij}^\alpha + \varphi_{\alpha\beta} \bar{x}_i^\alpha \bar{x}_j^\beta \\ &= \varphi_\alpha \bar{\nu}^\alpha \bar{h}_{ij} + \varphi_{\alpha\beta} \bar{x}_i^\alpha \bar{x}_j^\beta, \end{aligned}$$

where

$$(5.36) \quad \bar{x} = (\tau, x^i, \dots, x^n), \quad \tau = \text{const},$$

is the embedding of the coordinate slice, and we conclude

$$(5.37) \quad \varphi_{\alpha\beta} \bar{x}_i^\alpha \bar{x}_j^\beta = -\varphi_\alpha \bar{\nu}^\alpha \bar{h}_{ij} = -\bar{h}_{ij}$$

as well as

$$(5.38) \quad \varphi_{\alpha\beta} \bar{\nu}^\beta = 0.$$

Differentiating (5.37) covariantly with respect to the induced metric of the coordinate slices, we infer

$$(5.39) \quad \varphi_{\alpha\beta\gamma} \bar{x}_i^\alpha \bar{x}_j^\beta \bar{x}_k^\gamma = -\bar{h}_{ij;k} = 0,$$

in view of (5.31), where we also used (5.38).

Finally, differentiating (5.34) covariantly, we conclude

$$(5.40) \quad 0 = \varphi_{\alpha\beta\gamma}\varphi^\alpha + \varphi_{\alpha\beta}\varphi_\gamma^\alpha.$$

We can now evaluate the term

$$(5.41) \quad -\eta_{\alpha\beta\gamma}\nu^\alpha x_i^\beta x_j^\gamma = -\varphi_{\alpha\beta\gamma}\nu^\alpha x_i^\beta x_j^\gamma$$

with the help of the relations (5.39) and (5.40) yielding

$$(5.42) \quad \begin{aligned} -\varphi_{\alpha\beta\gamma}\nu^\alpha x_i^\beta x_j^\gamma &= -\varphi_{0ij}\nu^0 + \tilde{\nu}\varphi_{k0j}\check{u}^k u_i + \tilde{\nu}\varphi_{ki0}\check{u}^k u_j \\ &= -\tilde{\nu}\bar{h}_i^k \bar{h}_{kj} - \tilde{\nu}\bar{h}_k^m \bar{h}_{mi}\check{u}^k u_j - \tilde{\nu}\bar{h}_k^m \bar{h}_{mj}\check{u}^k u_i \\ &\quad + K_N \tilde{\nu} u_i u_j, \end{aligned}$$

where we applied the Ricci identities.

The indices of \bar{h}_{ij} are raised with the help of the metric \bar{g}_{ij} .

Taking then (5.31) into account completes the proof of the lemma. \square

The equation (5.27) can be even simplified further by using the same argument as in the case of its Riemannian analogue, cf. [4, Lemma 5.8].

Let $\eta = \eta(\tau)$ be a positive solution of the ordinary differential equation

$$(5.43) \quad \dot{\eta} = -\frac{\bar{H}}{n}\eta = -\bar{\kappa}\eta,$$

notice that η is defined for all $\tau \in \mathbb{R}$, and set

$$(5.44) \quad \chi = \tilde{v}\eta.$$

Then we can prove

5.7. Lemma. *The function χ satisfies the evolution equation*

$$(5.45) \quad \dot{\chi} - \dot{\Phi}F^{ij}\chi_{ij} = -\dot{\Phi}F^{ij}h_{ki}h_j^k\chi + [(\Phi - \tilde{f}) + \dot{\Phi}F]v\bar{\kappa}\chi + f_\alpha x_i^\alpha u^i v\chi,$$

for any value of K_N .

Proof. Differentiating (5.44) we deduce

$$(5.46) \quad \begin{aligned} \dot{\chi} - \dot{\Phi}F^{ij}\chi_{ij} &= \{\dot{\tilde{v}} - \dot{\Phi}F^{ij}\tilde{v}_{ij}\}\eta + \{\dot{u} - \dot{\Phi}F^{ij}u_{ij}\}\tilde{v}\eta \\ &\quad - 2\dot{\eta}\dot{\Phi}F^{ij}\tilde{v}_i u_j - \tilde{v}\dot{\eta}\dot{\Phi}F^{ij}u_i u_j. \end{aligned}$$

Now we first observe

$$(5.47) \quad \begin{aligned} \dot{\eta} &= -\frac{\dot{H}}{n}\eta - \frac{\bar{H}}{n}\dot{\eta} = -\frac{1}{n}\{|\bar{A}|^2 + \bar{R}_{\alpha\beta}\bar{\nu}^\alpha\bar{\nu}^\beta\}\eta + \bar{\kappa}^2\eta \\ &= K_N\eta. \end{aligned}$$

Secondly, from

$$(5.48) \quad \tilde{v}^2 = 1 + \|Du\|^2$$

we derive

$$(5.49) \quad \begin{aligned} \tilde{v}_i &= v u_{ij} u^j = -h_{ij}u^j + \bar{h}_{ij}u^j v \\ &= -h_{ij}u^j + \bar{\kappa}u_i \tilde{v}, \end{aligned}$$

hence we obtain

$$(5.50) \quad -2\dot{\eta}\dot{\Phi}F^{ij}\tilde{v}_i u_j = -2\bar{\kappa}\eta\dot{\Phi}F^{ij}h_{ik}u^k u_j + 2\bar{\kappa}^2\dot{F}^{ij}u_i u_j \tilde{v}\eta.$$

Inserting (5.47) and (5.50) in (5.46) we conclude

$$(5.51) \quad \begin{aligned} \dot{\chi} - \dot{\Phi}F^{ij}\chi_{ij} &= \{\dot{\tilde{v}} - \dot{\Phi}F^{ij}\tilde{v}_{ij}\}\eta - \bar{\kappa}\{\dot{u} - \dot{\Phi}F^{ij}u_{ij}\}\tilde{v}\eta \\ &\quad - 2\bar{\kappa}\dot{\Phi}F^{ij}h_{ik}u^k u_j \eta + 2\bar{\kappa}^2\dot{\Phi}F^{ij}u_i u_j \tilde{v}\eta \\ &\quad - K_N\dot{\Phi}F^{ij}u_i u_j \tilde{v}\eta, \end{aligned}$$

from which the result immediately follows, in view of (5.24) and (5.27). \square

5.8. Remark. Since the flow stays in a compact subset and the hypersurfaces $M(t)$ are uniformly convex, there exist positive constants c_1, c_2 such that

$$(5.52) \quad 0 < c_1 \leq \chi \leq c_2.$$

This follows immediately from the observation that in a point, where $D\chi = 0$, there holds

$$(5.53) \quad h_{ij}u^j = 0,$$

hence $Du = 0$, and thus $\chi = \eta$.

In case of a Lorentzian space form the evolution equation for the second fundamental form is a rather simple expression.

5.9. Lemma. *The second fundamental form (h_i^j) satisfies the differential equation*

$$\begin{aligned}
(5.54) \quad \dot{h}_i^j - \dot{\Phi} F^{kl} h_{i,kl}^j &= -\dot{\Phi} F^{kl} h_{rk} h_l^r h_i^j + \dot{\Phi} F h_{ri} h^{rj} - (\Phi - \tilde{f}) h_i^k h_k^j \\
&\quad - \tilde{f}_{\alpha\beta} x_i^\alpha x_k^\beta g^{kj} - \tilde{f}_\alpha \nu^\alpha h_i^j + \dot{\Phi} F^{kl,rs} h_{kl;i} h_{rs}^j \\
&\quad + \ddot{\Phi} F_i F^j \\
&\quad + K_N \{ (\Phi - \tilde{f}) \delta_i^j + \dot{\Phi} F \delta_i^j - \dot{\Phi} F^{kl} g_{kl} h_i^j \}.
\end{aligned}$$

Proof. The evolution equation for h_i^j in a semi-Riemannian manifold has been derived in [6, Lemma 3.5].

If the ambient space is a Lorentzian space form N with curvature K_N , and if M is a spacelike hypersurface with normal ν , then the former equation reduces to (5.54).

For Riemannian space forms, this equation has already been established in [4, Corollary 5.4]. Of course, in the stationary case, this identity was first proved by J. Simons in [17]. \square

6. CURVATURE ESTIMATES

We are now able to prove the a priori estimate for the principal curvatures of the $M(t)$.

6.1. Lemma. *Consider the flow in a maximal interval $I = [0, T^*)$, choose $\Phi = \log$, and assume that the initial hypersurface M_2 is of class $C^{4,\alpha}$, where $F \in (K^*)$ and $0 < f \in C^{2,\alpha}(\bar{\Omega})$. Then there are positive constants k_1, k_2 , depending only on F, f and Ω , such that the principal curvatures κ_i are estimated by*

$$(6.1) \quad 0 < k_1 \leq \kappa_i \leq k_2.$$

Proof. It suffices to prove an upper estimate for κ_i , since $F|_{\partial T_+} = 0$.

We observe that u, \tilde{v} and χ are already uniformly bounded, and that χ is also uniformly positive, cf. Remark 5.8.

Let φ and w be defined respectively by

$$(6.2) \quad \varphi = \sup \{ h_{ij} \eta^i \eta^j : \|\eta\| = 1 \},$$

$$(6.3) \quad w = \log \varphi + \lambda \chi,$$

where λ is a large positive parameter to be specified later. We claim that w is bounded for a suitable choice of λ .

Let $0 < T < T^*$, and $x_0 = x_0(t_0)$, with $0 < t_0 \leq T$, be a point in $M(t_0)$ such that

$$(6.4) \quad \sup_{M_0} w < \sup\{ \sup_{M(t)} w : 0 < t \leq T \} = w(x_0).$$

We then introduce a Riemannian normal coordinate system (ξ^i) at $x_0 \in M(t_0)$ such that at $x_0 = x(t_0, \xi_0)$ we have

$$(6.5) \quad g_{ij} = \delta_{ij} \quad \text{and} \quad \varphi = h_n^n.$$

Let $\tilde{\eta} = (\tilde{\eta}^i)$ be the contravariant vector field defined by

$$(6.6) \quad \tilde{\eta} = (0, \dots, 0, 1),$$

and set

$$(6.7) \quad \tilde{\varphi} = \frac{h_{ij}\tilde{\eta}^i\tilde{\eta}^j}{g_{ij}\tilde{\eta}^i\tilde{\eta}^j}.$$

$\tilde{\varphi}$ is well defined in neighbourhood of (t_0, ξ_0) .

Now, define \tilde{w} by replacing φ by $\tilde{\varphi}$ in (6.3); then, \tilde{w} assumes its maximum at (t_0, ξ_0) . Moreover, at (t_0, ξ_0) we have

$$(6.8) \quad \dot{\tilde{\varphi}} = \dot{h}_n^n,$$

and the spatial derivatives do also coincide; in short, at (t_0, ξ_0) $\tilde{\varphi}$ satisfies the same differential equation (5.54) as h_n^n . For the sake of greater clarity, let us therefore treat h_n^n like a scalar and pretend that w is defined by

$$(6.9) \quad w = \log h_n^n + \lambda\chi.$$

At (t_0, ξ_0) we have $\dot{w} \geq 0$, and, in view of the maximum principle, we deduce from (5.5), (5.54), and (5.45)

$$(6.10) \quad \begin{aligned} 0 \leq & \dot{\Phi}Fh_n^n - (\Phi - \tilde{f})h_n^n + \lambda c_1 - \lambda\epsilon_0\dot{\Phi}FH\chi \\ & + \lambda c_1[(\Phi - \tilde{f}) + \dot{\Phi}F] \\ & + \dot{\Phi}F^{ij}(\log h_n^n)_i(\log h_n^n)_j \\ & + \{\ddot{\Phi}F_nF^n + \dot{\Phi}F^{kl,rs}h_{kl;n}h_{rs;n}\}(h_n^n)^{-1}, \end{aligned}$$

where we have estimated bounded terms by a constant c_1 , assumed that h_n^n, λ are larger than 1, and used (5.5) as well as the simple observation

$$(6.11) \quad |F^{ij}h_j^k\eta_{ik}| \leq \|\eta\|F$$

valid for any tensor field (η_{ik}) .

Now, the last term in (6.10) is estimated from above by

$$(6.12) \quad \{\ddot{\Phi}F_nF^n + \dot{\Phi}F^{-1}F_nF^n\}(h_n^n)^{-1} - \dot{\Phi}F^{ij}h_{in;n}h_{jn;n}(h_n^n)^{-2},$$

cf. [6, Lemma 1.5], where the sum in the braces vanishes, due to the choice of Φ . Moreover, because of the Codazzi equation, we have

$$(6.13) \quad h_{in;n} = h_{nn;i},$$

and hence, we conclude that (6.12) is bounded from above by

$$(6.14) \quad - (h_n^n)^{-2}\dot{\Phi}F^{ij}h_{n;i}^nh_{n;j}^n.$$

Thus, the terms in (6.10) containing the derivatives of h_n^n sum up to something non-positive.

Choosing then in (6.10) λ such that

$$(6.15) \quad 2 \leq \lambda \epsilon_0 \chi$$

we derive

$$(6.16) \quad \begin{aligned} 0 \leq & -\dot{\Phi}FH - (\Phi - \tilde{f})h_n^n \\ & + \lambda c_1[(\Phi - \tilde{f}) + \dot{\Phi}F] + \lambda c_1. \end{aligned}$$

We now observe that $\dot{\Phi}F = 1$, and deduce in view of (5.23) that h_n^n is a priori bounded at (t_0, ξ_0) . \square

The result of the preceding lemma can be restated as a uniform estimate for the functions $u(t) \in C^2(S^n)$. Since, moreover, the principal curvatures of the flow hypersurfaces are not only bounded, but also uniformly bounded away from zero, in view of (5.23) and the assumption that F vanishes on $\partial\Gamma_+$, we conclude that F is uniformly elliptic on $M(t)$.

7. CONVERGENCE TO A STATIONARY SOLUTION

We are now ready to prove Theorem 0.1. Let $M(t)$ be the flow with initial hypersurface $M_0 = M_2$. Let us look at the scalar version of the flow

$$(7.1) \quad \frac{\partial u}{\partial t} = -v(\Phi - \tilde{f}),$$

cf. [6, equ. (3.5)].

This is a scalar parabolic differential equation defined on the cylinder

$$(7.2) \quad Q_{T^*} = [0, T^*) \times S^n$$

with initial value $u(0) = u_2 \in C^{4,\alpha}(S^n)$. In view of the a priori estimates, which we have established in the preceding sections, we know that

$$(7.3) \quad |u|_{2,0,S^n} \leq c$$

and

$$(7.4) \quad \Phi(F) \text{ is uniformly elliptic in } u$$

independent of t . Moreover, $\Phi(F)$ is concave, and thus, we can apply the regularity results of [10, Chapter 5.5] to conclude that uniform $C^{2,\alpha}$ -estimates are valid, leading further to uniform $C^{4,\alpha}$ -estimates due to the regularity results for linear operators.

Therefore, the maximal time interval is unbounded, i.e. $T^* = \infty$.

Now, integrating (7.1) with respect to t , and observing that the right-hand side is non-positive, yields

$$(7.5) \quad u(0, x) - u(t, x) = \int_0^t v(\Phi - \tilde{f}) \geq c \int_0^t (\Phi - \tilde{f}),$$

i.e.,

$$(7.6) \quad \int_0^\infty |\Phi - \tilde{f}| < \infty \quad \forall x \in S^n$$

Hence, for any $x \in S^n$ there is a sequence $t_k \rightarrow \infty$ such that $(\Phi - \tilde{f}) \rightarrow 0$.

On the other hand, $u(\cdot, x)$ is monotone decreasing and therefore

$$(7.7) \quad \lim_{t \rightarrow \infty} u(t, x) = \tilde{u}(x)$$

exists and is of class $C^{4,\alpha}(S^n)$ in view of the a priori estimates. We, finally, conclude that \tilde{u} is a stationary solution of our problem, and that

$$(7.8) \quad \lim_{t \rightarrow \infty} (\Phi - \tilde{f}) = 0.$$

REFERENCES

- [1] A.D. Alexandrov, *On the theory of mixed volumes of convex bodies*, Mat. Sb. **3** (1938), 27–46.
- [2] Shiu Yuen Cheng and Shing Tung Yau, *On the regularity of the solution of the n -dimensional Minkowski problem*, Comm. Pure Appl. Math. **29** (1976), no. 5, 495–516.
- [3] Claus Gerhardt, *Closed Weingarten hypersurfaces in Riemannian manifolds*, J. Diff. Geom. **43** (1996), 612–641, pdf file.
- [4] ———, *Closed Weingarten hypersurfaces in space forms*, Geometric Analysis and the Calculus of Variations (J Jost, ed.), International Press, Boston, 1996, pdf file, p. 71–98.
- [5] ———, *Hypersurfaces of prescribed Weingarten curvature*, Math. Z. **224** (1997), 167–194, download pdf file.
- [6] ———, *Hypersurfaces of prescribed curvature in Lorentzian manifolds*, Indiana Univ. Math. J. **49** (2000), 1125–1153, arXiv:math.DG/0409457.
- [7] ———, *Analysis II*, International Press, Somerville, MA, 2005, 404 pp.
- [8] ———, *Minkowski type problems for convex hypersurfaces in the sphere*, 2005, arXiv:math.DG/0509217, 25 pages.
- [9] Bo Guan and Pengfei Guan, *Convex hypersurfaces of prescribed curvatures.*, Ann. Math. **156** (2002), 655–673.
- [10] N. V. Krylov, *Nonlinear elliptic and parabolic equations of the second order*, Mathematics and its Applications (Soviet Series), vol. 7, D. Reidel Publishing Co., Dordrecht, 1987.
- [11] Hans Lewy, *On differential geometry in the large. I. Minkowski's problem*, Trans. Amer. Math. Soc. **43** (1938), no. 2, 258–270.
- [12] H. Minkowski, *Volumen und Oberfläche.*, Math. Annalen **43** (1903), 447–495.
- [13] Louis Nirenberg, *The Weyl and Minkowski problems in differential geometry in the large*, Comm. Pure Appl. Math. **6** (1953), 337–394.
- [14] Barrett O'Neill, *Semi-Riemannian geometry. With applications to relativity.*, Pure and Applied Mathematics, 103. New York-London etc.: Academic Press. XIII, 1983.
- [15] A. V. Pogorelov, *Regularity of a convex surface with given Gaussian curvature*, Mat. Sbornik N.S. **31(73)** (1952), 88–103.
- [16] Richard Sacksteder, *On hypersurfaces with no negative sectional curvatures*, Amer. J. Math. **82** (1960), 609–630.
- [17] James Simons, *Minimal varieties in riemannian manifolds*, Ann. of Math. (2) **88** (1968), 62–105.

RUPRECHT-KARLS-UNIVERSITÄT, INSTITUT FÜR ANGEWANDTE MATHEMATIK, IM NEUEN-
HEIMER FELD 294, 69120 HEIDELBERG, GERMANY

E-mail address: `gerhardt@math.uni-heidelberg.de`

URL: `http://www.math.uni-heidelberg.de/studinfo/gerhardt/`