

**DERIVING A COMPLETE SET OF EIGENDISTRIBUTIONS
FOR A GRAVITATIONAL WAVE EQUATION DESCRIBING
THE QUANTIZED INTERACTION OF GRAVITY WITH A
YANG-MILLS FIELD IN CASE THE CAUCHY
HYPERSURFACE IS NON-COMPACT**

CLAUS GERHARDT

ABSTRACT. In a recent paper we quantized the interaction of gravity with a Yang-Mills and Higgs field and obtained as a result a gravitational wave equation in a globally hyperbolic spacetime. Assuming that the Cauchy hypersurfaces are compact we proved a spectral resolution for the wave equation by applying the method of separation of variables. In this paper we extend the results to the case when the Cauchy hypersurfaces are non-compact by considering a Gelfand triplet and applying the nuclear spectral theorem.

CONTENTS

| | |
|--|----|
| 1. Introduction | 1 |
| 2. The nuclear spectral theorem | 4 |
| 3. The eigendistributions are smooth functions | 8 |
| 4. The positivity of the eigenvalues | 11 |
| References | 15 |

1. INTRODUCTION

In a recent paper [3] we quantized the interaction of gravity with a Yang-Mills and Higgs field and obtained as a result a gravitational wave equation of the form

$$(1.1) \quad \frac{1}{32} \frac{n^2}{n-1} \ddot{u} - (n-1)t^{2-\frac{4}{n}} \Delta u - \frac{n}{2} t^{2-\frac{4}{n}} R u + \alpha_1 \frac{n}{8} t^{2-\frac{4}{n}} F_{ij} F^{ij} u + \alpha_2 \frac{n}{4} t^{2-\frac{4}{n}} \gamma_{ab} \sigma^{ij} \Phi_i^a \Phi_i^b u + \alpha_2 \frac{n}{2} m t^{2-\frac{4}{n}} V(\Phi) u + n t^2 \Lambda u = 0,$$

Date: May 13, 2016.

2000 Mathematics Subject Classification. 83,83C,83C45.

Key words and phrases. unified field theory, quantization of gravity, quantum gravity, Yang-Mills fields, eigendistributions, Gelfand triple, nuclear spectral theorem, mass gap.

in a globally hyperbolic spacetime

$$(1.2) \quad Q = (0, \infty) \times \mathcal{S}_0$$

describing the interaction of a given complete Riemannian metric σ_{ij} in \mathcal{S}_0 with a given Yang-Mills and Higgs field; R is the scalar curvature of σ_{ij} , V is the potential of the Higgs field, m a positive constant, α_1, α_2 are positive coupling constants and the other symbols should be self-evident. The existence of the time variable, and its range, is due to the quantization process.

1.1. Remark. For the results and arguments in that paper it was completely irrelevant that the values of the Higgs field Φ lie in a Lie algebra, i.e., Φ could also be just an arbitrary scalar field, or we could consider a Higgs field as well as an another arbitrary scalar field. Hence, let us stipulate that the Higgs field could also be just an arbitrary scalar field.

If \mathcal{S}_0 is compact we also proved a spectral resolution of equation (1.1) by first considering a stationary version of the hyperbolic equation, namely, the elliptic eigenvalue equation

$$(1.3) \quad \begin{aligned} & - (n-1)\Delta v - \frac{n}{2}Rv + \alpha_1 \frac{n}{8} F_{ij} F^{ij} v \\ & + \alpha_2 \frac{n}{4} \gamma_{ab} \sigma^{ij} \Phi_i^a \Phi_j^b v + \alpha_2 \frac{n}{2} m V(\Phi) v = \mu v. \end{aligned}$$

It has countably many solutions (v_i, μ_i) such that

$$(1.4) \quad \mu_0 < \mu_1 \leq \mu_2 \leq \dots,$$

$$(1.5) \quad \lim \mu_i = \infty.$$

Let v be an eigenfunction with eigenvalue $\mu > 0$, then we looked at solutions of (1.1) of the form

$$(1.6) \quad u(x, t) = w(t)v(x).$$

u is then a solution of (1.1) provided w satisfies the implicit eigenvalue equation

$$(1.7) \quad -\frac{1}{32} \frac{n^2}{n-1} \ddot{w} - \mu t^{2-\frac{4}{n}} w - nt^2 \Lambda w = 0,$$

where Λ is the eigenvalue.

This eigenvalue problem we also considered in a previous paper and proved that it has countably many solutions (w_i, Λ_i) with finite energy, i.e.,

$$(1.8) \quad \int_0^\infty \{|\dot{w}_i|^2 + (1+t^2 + \mu t^{2-\frac{4}{n}})|w_i|^2\} < \infty.$$

More precisely, we proved, cf. [2, Theorem 6.7],

1.2. Theorem. *Assume $n \geq 2$ and \mathcal{S}_0 to be compact and let (v, μ) be a solution of the eigenvalue problem (1.3) with $\mu > 0$, then there exist countably many solutions (w_i, Λ_i) of the implicit eigenvalue problem (1.7) such that*

$$(1.9) \quad \Lambda_i < \Lambda_{i+1} < \cdots < 0,$$

$$(1.10) \quad \lim_i \Lambda_i = 0,$$

and such that the functions

$$(1.11) \quad u_i = w_i v$$

are solutions of the wave equation (1.1). The transformed eigenfunctions

$$(1.12) \quad \tilde{w}_i(t) = w_i(\lambda_i^{\frac{n}{4(n-1)}} t),$$

where

$$(1.13) \quad \lambda_i = (-\Lambda_i)^{-\frac{n-1}{n}},$$

form a basis of $L^2(\mathbb{R}_+^*, \mathbb{C})$ and also of the Hilbert space H defined as the completion of $C_c^\infty(\mathbb{R}_+^*, \mathbb{C})$ under the norm of the scalar product

$$(1.14) \quad \langle w, \tilde{w} \rangle_1 = \int_0^\infty \{\bar{w}' \tilde{w}' + t^2 \bar{w} \tilde{w}\},$$

where a prime or a dot denotes differentiation with respect to t .

In this paper we want to extend this spectral resolution to the case when \mathcal{S}_0 is non-compact. Denote by A the elliptic differential operator on the left-hand side of (1.3), then, assuming that its coefficients are smooth with bounded C^m -norms for any $m \in \mathbb{N}$, we have a self-adjoint operator in $H = L^2(\mathcal{S}_0)$ and a Gelfand triplet

$$(1.15) \quad \mathcal{S} \subset H \subset \mathcal{S}'$$

such that we can apply the nuclear spectral theorem of Gelfand-Maurin leading to a complete set of eigendistributions

$$(1.16) \quad f(\lambda) \in \mathcal{S}', \quad \lambda \in \Lambda,$$

of A , where Λ is a measure space. For almost every $\lambda \in \Lambda$ we have $0 \neq f(\lambda)$ and $f(\lambda)$ is a solution of the eigenvalue equation

$$(1.17) \quad Af(\lambda) = a(\lambda)f(\lambda)$$

where

$$(1.18) \quad a : \Lambda \rightarrow \sigma(A)$$

is a measurable function having $\sigma(A)$ as its essential range. Since the $f(\lambda)$ are distributions and A is uniformly elliptic and smooth, the $f(\lambda)$ are also smooth, and since they are also tempered distributions we could prove that the eigenvalues satisfy

$$(1.19) \quad a(\lambda) > 0$$

for a.e. λ . Hence, the separation of variables, described in (1.6), can be applied with an eigenfunction v be replaced by an eigendistribution f . Since all eigenvalues $a(\lambda)$ are strictly positive this can be considered to be a spectral resolution of the wave equation. The smooth functions

$$(1.20) \quad u_i = w_i f(\lambda)$$

are classical solutions of the wave equation (1.1) with bounded temporal energy and locally bounded spatial energy.

2. THE NUCLEAR SPECTRAL THEOREM

We assume that $(\mathcal{S}_0, \sigma_{ij})$ is complete and that there exists a compact subset $K_0 \subset \mathcal{S}_0$ and a chart (U_0, x) such that

$$(2.1) \quad \mathcal{S}_0 \setminus K_0 \subset U_0$$

and

$$(2.2) \quad \Omega_0 = x(U_0) = \mathbb{R}^n \setminus \bar{B}_{R_0}(0).$$

Moreover, we require

2.1. Assumption. (i) The metric σ_{ij} and the lower order coefficients of the elliptic operator on the left-hand side of equation of the equation (1.3) on page 2 are smooth with bounded C^m -norms for any $m \in \mathbb{N}$. We call the elliptic operator A .

(ii) The metric σ_{ij} is uniformly elliptic.

The last assumption implies that the radial distance from a center $x_0 \in K_0$,

$$(2.3) \quad r(x) = d(x, x_0),$$

and the Euclidean distance $|x|$ are equivalent in Ω_0 , i.e., there are constants c_1, c_2 such that

$$(2.4) \quad r(x) \leq c_1|x| \leq c_2r(x) \quad \forall x \in \Omega_0$$

and hence the Schwartz space of rapidly decreasing test functions in \mathcal{S}_0 can be identified with the Schwartz space in \mathbb{R}^n . We shall denote the Schwartz space by

$$(2.5) \quad \mathcal{S} = \mathcal{S}(\mathcal{S}_0)$$

and its dual space, the tempered distributions, by

$$(2.6) \quad \mathcal{S}' = \mathcal{S}'(\mathcal{S}_0).$$

The topology of \mathcal{S} is defined by a sequence of norms

$$(2.7) \quad |\varphi|_{m,k} = \sup_{x \in \mathcal{S}_0} (1 + r(x)^2)^k \sum_{|\alpha| \leq m} |D^\alpha \varphi(x)|.$$

\mathcal{S} is a Fréchet space and also a nuclear space, cf. [8, Example 5, p. 107]. The differential operator A defined by the left-hand side of (1.3) on page 2 is a

continuous map from \mathcal{S} into \mathcal{S} in view of Assumption 2.1. A is a self-adjoint linear operator in $L^2(\mathcal{S}_0, \mathbb{C})$ and

$$(2.8) \quad \mathcal{S} \subset D(A)$$

a dense subspace. By duality A can also be defined on the dual space \mathcal{S}' , namely, let $f \in \mathcal{S}'$, then

$$(2.9) \quad \langle Af, \varphi \rangle = \langle f, A\varphi \rangle \quad \forall \varphi \in \mathcal{S},$$

where the self-adjointness of A has been used.

2.2. Definition. $f \in \mathcal{S}'$ is said to be an eigendistribution of A with eigenvalue $\mu \in \mathbb{R}$, i.e.,

$$(2.10) \quad Af = \mu f,$$

iff

$$(2.11) \quad \langle Af, \varphi \rangle = \mu \langle f, \varphi \rangle \quad \forall \varphi \in \mathcal{S},$$

or equivalently, iff

$$(2.12) \quad \langle f, A\varphi \rangle = \mu \langle f, \varphi \rangle \quad \forall \varphi \in \mathcal{S}.$$

A setting where we have a self-adjoint operator A in a separable Hilbert space H , a dense subspace

$$(2.13) \quad E \subset H$$

which is also a nuclear space (in a finer topology) with dual space E' such that

$$(2.14) \quad E \subset H \subset E',$$

where the imbedding of E into H is continuous, a property which we already specified by speaking of a *finer* topology, and where, moreover,

$$(2.15) \quad A : E \rightarrow E$$

is continuous, is known as a *rigged Hilbert space* setting, though, usually, E' is replaced by the space of antilinear functionals. However, since we do not use Dirac's ket notation, we shall consider E' .

In such a framework a nuclear spectral theorem has been proved by Gelfand and Maurin, cf. [1, Theorem 5', p. 126], [5, Satz 2] and [6, Chap. XVIII, p. 333] which we shall formulate and prove for a single self-adjoint operator A and not for a family of strongly commuting operators. The proof closely follows the one given by Maurin in [5, Satz 2]. Since this paper is written in German we like to include a proof for the convenience of the reader.

2.3. Theorem (Maurin). *Let H be a separable complex Hilbert space, A a densely defined self-adjoint operator, $E \subset H$ a dense subspace which also carries topology such that it is a nuclear space and assume that the imbedding*

in (2.13) and the map A in (2.15) are continuous, then there exists a locally compact measure space Λ , a finite positive measure μ , a measurable function

$$(2.16) \quad a : \Lambda \rightarrow \sigma(A) \subset \mathbb{R},$$

and a unitary operator

$$(2.17) \quad U : H \rightarrow L^2(\Lambda, \mathbb{C}, \mu)$$

such that, if we set

$$(2.18) \quad \hat{u} = Uu, \quad \forall u \in H,$$

$$(2.19) \quad \hat{A} = UAU^{-1},$$

we have

$$(2.20) \quad u \in D(A) \iff a\hat{u} \in L^2(\Lambda, \mu),$$

$$(2.21) \quad \hat{A}\hat{u} = a\hat{u} \quad \forall u \in D(A)$$

and for μ a.e. $\lambda \in \Lambda$ the mapping

$$(2.22) \quad f(\lambda) : \varphi \in E \rightarrow \hat{\varphi}(\lambda) \in \mathbb{C}$$

is continuous in E and does not vanish identically, i.e.,

$$(2.23) \quad 0 \neq f(\lambda) \in E'$$

and hence

$$(2.24) \quad \hat{\varphi}(\lambda) = \langle f(\lambda), \varphi \rangle \quad \forall \varphi \in E.$$

Moreover, (2.21) implies

$$(2.25) \quad \langle f(\lambda), A\varphi \rangle = \hat{A}\hat{\varphi}(\lambda) = a(\lambda)\hat{\varphi}(\lambda) = a(\lambda)\langle f(\lambda), \varphi \rangle \quad \forall \varphi \in E$$

for a.e. $\lambda \in \Lambda$, or equivalently,

$$(2.26) \quad Af(\lambda) = a(\lambda)f(\lambda) \quad \text{for a.e. } \lambda \in \Lambda.$$

The generalized eigenvectors $f(\lambda)$ are complete, since

$$(2.27) \quad \|\varphi\|^2 = \|\hat{\varphi}\|^2 = \int_{\Lambda} |\hat{\varphi}(\lambda)|^2 d\mu \quad \forall \varphi \in E,$$

and hence,

$$(2.28) \quad \hat{\varphi}(\lambda) = 0 \quad \text{for a.e. } \lambda \in \Lambda,$$

is equivalent to $\varphi = 0$.

Proof. The first part of the theorem is due to the multiplicative form of the spectral theorem, cf. [7, Theorem VIII.4, p. 260]. Let us remark that we used a different version of von Neumann's spectral theorem than Maurin which simplifies the proof slightly, especially the completeness part. Note that the spectrum

$$(2.29) \quad \sigma(A) = \sigma(\hat{A})$$

is the essential range of a .

To prove (2.22) and the following claims, we observe that the imbedding

$$(2.30) \quad j : E \rightarrow H$$

is continuous and therefore also nuclear, hence there is a semi-norm $\|\cdot\|_p$ on E sequences $u_k \in H$, $f_k \in E'$ such that

$$(2.31) \quad j(\varphi) = \sum_k \langle f_k, \varphi \rangle u_k \quad \forall \varphi \in E$$

and

$$(2.32) \quad \sum_k \|f_k\|_{-p} \|u_k\| = \sum_k \|f_k\|_{-p} \|\hat{u}_k\| < \infty,$$

where $\|\cdot\|_{-p}$ is the dual norm in E'

$$(2.33) \quad \|f_k\|_{-p} = \sup_{\|\varphi\|_p=1} |\langle f_k, \varphi \rangle|.$$

We shall show that the mapping in (2.30), which, when composed with U , can now be expressed as

$$(2.34) \quad \varphi \rightarrow \hat{\varphi}(\lambda) = \sum_k \langle f_k, \varphi \rangle \hat{u}_k(\lambda)$$

is continuous in E and not identically 0 for a.e. $\lambda \in \Lambda$.

Indeed, without loss of generality we may assume

$$(2.35) \quad \|u_k\| = 1$$

to deduce from (2.32)

$$(2.36) \quad \begin{aligned} \sum_k \|f_k\|_{-p} &= \sum_k \|f_k\|_{-p} \|u_k\| = \sum_k \|f_k\|_{-p} \|u_k\|^2 \\ &= \sum_k \|f_k\|_{-p} \int_{\Lambda} |\hat{u}_k(\lambda)|^2 = \int_{\Lambda} \sum_k \|f_k\|_{-p} |\hat{u}_k(\lambda)|^2 < \infty, \end{aligned}$$

hence

$$(2.37) \quad \sum_k \|f_k\|_{-p} |\hat{u}_k(\lambda)|^2 \equiv c_1(\lambda)^2 < \infty$$

for a.e. $\lambda \in \Lambda$, and

$$(2.38) \quad c_1(\cdot) \in L^2(\Lambda, \mu).$$

To prove (2.22) we now estimate

$$\begin{aligned}
|\hat{\varphi}(\lambda)|^2 &= \left| \sum_k \langle f_k, \varphi \rangle \hat{u}_k(\lambda) \right|^2 \leq \left(\sum_k \|f_k\|_{-p} \|\varphi\|_p |\hat{u}_k(\lambda)| \right)^2 \\
&= \left(\sum_k \|f_k\|_{-p}^{\frac{1}{2}} \|\varphi\|_p \|f_k\|_{-p}^{\frac{1}{2}} |\hat{u}_k(\lambda)| \right)^2 \\
(2.39) \quad &\leq \left(\sum_k \|f_k\|_{-p} \|\varphi\|_p^2 \right) \left(\sum_k \|f_k\|_{-p} |\hat{u}_k(\lambda)|^2 \right) \\
&= c_1(\lambda)^2 \sum_k \|f_k\|_{-p} \|\varphi\|_p^2 < \infty,
\end{aligned}$$

in view of (2.36) and (2.37).

The fact that the mapping in (2.34) does not vanish identically for a.e. $\lambda \in \Lambda$ is proved in the lemma below. This completes the proof of the theorem, since the other properties are evident. \square

2.4. Lemma. *The mapping (2.34) does not vanish identically in E for a.e. $\lambda \in \Lambda$.*

Proof. We argue by contradiction and assume that there exists a measurable set $\Lambda_0 \subset \Lambda$ with positive measure such that

$$(2.40) \quad \hat{\varphi}(\lambda) = 0 \quad \forall (\lambda, \varphi) \in \Lambda_0 \times E.$$

Let χ_0 be the characteristic function of Λ_0 and set

$$(2.41) \quad u_0 = U^{-1} \chi_0,$$

then

$$(2.42) \quad 0 \neq u_0 \in H.$$

Let $\varphi_k \in E$ be sequence converging to u_0 , then

$$(2.43) \quad \|u_0\|^2 = \lim_k \langle \varphi_k, u_0 \rangle = \lim_k \int_{\Lambda} \bar{\varphi}_k \chi_0 = 0,$$

a contradiction. \square

3. THE EIGENDISTRIBUTIONS ARE SMOOTH FUNCTIONS

In our case $E = \mathcal{S}$ and A is a uniformly elliptic linear differential operator with smooth coefficients. Hence, we can prove:

3.1. Theorem. *Let A satisfy the Assumption 2.1 on page 4, then the solutions $f(\lambda) \in \mathcal{S}'$ of the eigenvalue problem*

$$(3.1) \quad Af(\lambda) = \mu f(\lambda)$$

belong to $C^\infty(\mathcal{S}_0)$ and for each $m \in \mathbb{N}$ and $R > 0$ $f(\lambda)$ can be estimated by

$$(3.2) \quad |f(\lambda)|_{m, B_R(x_0)} \leq c_m R^N \|f(\lambda)\|_{-p},$$

where $\|\cdot\|_p$ is one of the defining norms in \mathcal{S} such that

$$(3.3) \quad \|f(\lambda)\|_{-p} = \sup_{\|\varphi\|_p=1} |\langle f(\lambda), \varphi \rangle|$$

and N depends on n , $\|\cdot\|_p$, A and \mathcal{S}_0 , while c_m depends on m , A the eigenvalue μ and on \mathcal{S}_0 . $B_R(x_0)$ is a geodesic ball of radius R for a fixed $x_0 \in K_0 \subset \mathcal{S}_0$.

Proof. First we note that we can absorb the right-hand side of the eigenvalue equation into the left-hand side and simply consider the equation

$$(3.4) \quad Af(\lambda) = 0.$$

Hence, it is well-known that the distributional solutions is smooth and equation (3.4) can be understood in the classical sense, see e.g., [4, Theorem 3.2, p.125].

The important estimate (3.2) is due to the fact that $f(\lambda)$ is a tempered distribution. Since $f(\lambda) \in \mathcal{S}'$ we have

$$(3.5) \quad |\langle f(\lambda), \varphi \rangle| \leq c \sup_{x \in \mathcal{S}_0} (1 + r(x)^2)^k \sum_{|\alpha| \leq m_0} |D^\alpha \varphi(x)| \equiv c \|\varphi\|_p$$

and the dual norm

$$(3.6) \quad \|f(\lambda)\|_{-p} = c.$$

To prove (3.2) we fix $m \in \mathbb{N}$ and assume that

$$(3.7) \quad |f(\lambda)|_{m, B_{R_1}(x_0)} \leq c_0,$$

for some sufficiently large radius R_1 such that we only have to prove the estimate in the domain

$$(3.8) \quad B_R(0) \setminus \bar{B}_{R_0}(0),$$

where we now consider Euclidean balls, cf. the assumptions in (2.1) and (2.2) on page 4. Hence we may consider equation (3.4) to be a uniformly elliptic equation in an exterior region of Euclidean space with smooth coefficients.

Let $R > R_0$, then we first prove a priori estimates for $f(\lambda)$ in small balls

$$(3.9) \quad B_\rho(y) \Subset B_{2R}(0) \setminus B_{R_0}(0),$$

where

$$(3.10) \quad 2\rho < \rho_0 \leq 1$$

and ρ_0 is fixed.

Let

$$(3.11) \quad H_0^{m,2}(\Omega), \quad m \in \mathbb{N},$$

be the usual Sobolev spaces, where

$$(3.12) \quad \Omega \subset \mathbb{R}^n$$

is an open set, to be defined as the completion of $C_c^\infty(\Omega)$ under the norm

$$(3.13) \quad \|\varphi\|_{m,2}^2 = \int_{\Omega} \sum_{|\alpha| \leq m} |D^\alpha \varphi|^2.$$

$H_0^{m,2}(\Omega)$ is a Hilbert space. Its dual space is denoted by

$$(3.14) \quad H^{-m,2}(\Omega)$$

and its elements are the distributions $f \in \mathcal{D}'(\Omega)$ which can be written in the form

$$(3.15) \quad f = \sum_{|\alpha| \leq m} D^\alpha u_\alpha,$$

where

$$(3.16) \quad u_\alpha \in L^2(\Omega)$$

and the dual norm of f is equal to

$$(3.17) \quad \|f\|_{-m,2} = \left(\sum_{|\alpha| \leq m} \|u_\alpha\|_2^2 \right)^{\frac{1}{2}}.$$

The Sobolev imbedding theorem states that

$$(3.18) \quad m > \frac{n}{2} \quad \Longrightarrow \quad H_0^{m,2}(\Omega) \hookrightarrow C^0(\Omega)$$

such that

$$(3.19) \quad |u|_0 \leq c \|u\|_{m,2} \quad \forall u \in H_0^{m,2}(\Omega),$$

where c only depends on m and n .

As a corollary we deduce

$$(3.20) \quad m > \frac{n}{2} \quad \Longrightarrow \quad H_0^{m+m_0,2}(\Omega) \hookrightarrow C^{m_0,0}(\Omega)$$

with a corresponding estimate

$$(3.21) \quad |u|_{m_0,0} \leq c \|u\|_{m+m_0,2},$$

where $c = c(n, m, m_0)$.

Hence, for any ball

$$(3.22) \quad B_{\rho_0}(y) \subset B_{2R}(0)$$

$f(\lambda)$ can be considered to belong to

$$(3.23) \quad f(\lambda) \in H^{-(m_0+n),2}(B_{\rho_0}(y))$$

with norm

$$(3.24) \quad \|f(\lambda)\|_{-(n+m_0),2} \leq c R^{2k}$$

in view of the estimate (3.5), where we also assume $R_0 > 1$; the constant c depends on n, m_0, k and the constant in (3.5).

From the proofs of [4, Theorem 3.1, p. 123] and [4, Theorem 3.2, p. 125] we then deduce that for any $m \in \mathbb{N}$ there exists $\rho < \rho_0$, ρ depending only on

the Lipschitz constant of the metric σ_{ij} , m , n and m_0 such that the C^m -norm of the solution $f(\lambda)$ of equation (3.4) can be estimated by

$$(3.25) \quad |f(\lambda)|_{m, B_\rho(y)} \leq c_\rho R^{2k},$$

where c_ρ also depends on the C^m -norms of the coefficients of A and on the ellipticity constants.

Now

$$(3.26) \quad (4R)^n 2^n \rho^{-n}$$

balls

$$(3.27) \quad B_\rho(y) \subset B_{2R}(0)$$

cover the closed ball $\bar{B}_R(0)$, hence we conclude

$$(3.28) \quad |f(\lambda)|_{m, B_R(0) \setminus K_0} \leq cR^{2k+n},$$

where $c = c(\rho, m, m_0, n, A)$. \square

4. THE POSITIVITY OF THE EIGENVALUES

To apply the separation of variables method to find a complete set of eigensolutions for the wave equation the eigenvalues of the elliptic operator have to be positive. In this section we shall prove that eigenvalues of the eigenvalue equation (3.1) on page 8 are always strictly positive provided some rather weak assumptions are satisfied.

Let us start with the following lemma:

4.1. Lemma. *Let A be the differential operator on the left-hand side of (1.3) on page 2 and let us write the operator in the form*

$$(4.1) \quad Av = -(n-1)\Delta v - \frac{n}{2}Rv + Gv + \alpha_2 \frac{n}{2}mV(\Phi)v,$$

where

$$(4.2) \quad 0 \leq G = \alpha_1 \frac{n}{8}F_{ij}F^{ij} + \alpha_2 \frac{n}{2}\gamma_{ab}\sigma^{ij}\Phi_i^a\Phi_j^b.$$

Assume there are positive constants ϵ_0 , δ , m_0 and R_1 such that

$$(4.3) \quad -\frac{n}{2}R + G + \alpha_2 \frac{n}{2}m_0V \geq \epsilon_0 r^{-2+\delta} \quad \forall x \notin B_{R_1}(x_0),$$

where $x_0 \in K_0$ is fixed and r is the geodesic distance to x_0 , then there exists $m_1 \geq m_0$ such that for all $m \geq m_1$ the quadratic form of A satisfies

$$(4.4) \quad \int_{B_{R_1}(x_0)} \|u\|^2 \leq \langle Au, u \rangle \quad \forall u \in H^{1,2}(\mathcal{S}_0),$$

provided

$$(4.5) \quad V(\Phi) > 0 \quad \text{a.e. in } \mathcal{S}_0.$$

Proof. The ball $B_{R_1}(x_0)$ is bounded, hence the imbedding of

$$(4.6) \quad H^{1,2}(B_{R_1}(x_0)) \hookrightarrow L^2(B_{R_1}(x_0))$$

is compact and we can apply a compactness lemma to conclude that for any $\epsilon > 0$ there is a constant c_ϵ such that

$$(4.7) \quad \int_{B_{R_1}(x_0)} |u|^2 \leq \epsilon \int_{B_{R_1}(x_0)} |Du|^2 + c_\epsilon \int_{B_{R_1}(x_0)} V(\Phi)|u|^2$$

for all $u \in H^{1,2}(B_{R_1}(x_0))$, in view of the assumption (4.5), cf. [3, Lemma 7.5].

Hence, we deduce

$$(4.8) \quad \int_{B_{R_1}(x_0)} |u|^2 \leq (n-1) \int_{B_{R_1}(x_0)} |Du|^2 + \int_{B_{R_1}(x_0)} \left(-\frac{n}{2}R + G\right) |u|^2 \\ + \alpha_2 \frac{n}{2} m \int_{B_{R_1}(x_0)} V(\Phi) |u|^2$$

for all $u \in H^{1,2}(B_{R_1}(x_0))$ provided m is sufficiently large

$$(4.9) \quad m \geq m_1.$$

Choosing $m_1 \geq m_0$ completes the proof of the lemma because of the assumption (4.3). \square

4.2. Theorem. *Under the assumptions of the preceding lemma and the general provisions in (2.1), (2.2) and Assumption 2.1 on page 4 the eigenvalue equation (3.1) on page 8 is only solvable if $\mu > 0$.*

Proof. Since the quadratic form of A is positive we immediately infer

$$(4.10) \quad \sigma(A) \subset \mathbb{R}_+,$$

hence the eigenvalue μ in (3.1) has to satisfy

$$(4.11) \quad 0 \leq \mu$$

so that we have to exclude the case

$$(4.12) \quad \mu = 0.$$

We argue by contradiction. Let

$$(4.13) \quad f \in \mathcal{S}' \cap C^\infty(\mathcal{S}_0)$$

be a solution of

$$(4.14) \quad Af = 0,$$

then we shall prove

$$(4.15) \quad f = 0.$$

Let $k \in \mathbb{N}$ and $R > R_1$ be large and let η be defined by

$$(4.16) \quad \eta(x) = \begin{cases} R^{-k}, & |x| \leq R, \\ |x|^{-k}, & |x| > R, \end{cases}$$

Then

$$(4.17) \quad f\eta^2 \in H^{1,2}(\mathcal{S}_0),$$

in view of the estimate (3.2) on page 8, which can be rephrased to

$$(4.18) \quad \sum_{|\alpha| \leq m} |D^\alpha f(x)| \leq c_m |x|^N \quad \forall |x| > R_0.$$

Here, we use the Euclidean distance.

Multiplying (4.14) by $f\eta^2$ and integrating by parts yields

$$(4.19) \quad \begin{aligned} 0 &\geq \int_{\mathcal{S}_0} \left\{ (n-1)|Df|^2\eta^2 - \frac{n}{2}R|f|^2 + G|f|^2 + \alpha_2 \frac{n}{2}m|f|^2 \right\} \eta^2 \\ &\quad - (n-1) \int_{\mathcal{S}_0 \setminus B_{R_1}(x_0)} 2|Df||f|\eta|D\eta| \\ &\geq \int_{B_R} \left\{ (n-1)|Df|^2 - \frac{n}{2}R|f|^2 + G|f|^2 + \alpha_2 \frac{n}{2}m|f|^2 \right\} R^{-2k} \\ &\quad + \int_{\mathcal{S}_0 \setminus B_R} \{ \epsilon_0 |x|^{-2+\delta} - c|x|^{-2} \} |f|^2 \eta^2, \end{aligned}$$

where c is a fixed constant depending only on the metric σ_{ij} and k .

The first integral is strictly positive unless f vanishes in $B_R(x_0)$, and the difference in the braces is also strictly positive if R is large enough. Hence we conclude

$$(4.20) \quad f \equiv 0. \quad \square$$

We can now prove a spectral resolution of the hyperbolic equation (1.1) on page 1 by choosing an eigendistribution $f = f(\lambda)$ with eigenvalue $\mu = a(\lambda)$ and look at solutions of (1.1) of the form

$$(4.21) \quad u(x, t) = w(t)f(x).$$

u is then a solution of (1.1) provided w satisfies the implicit eigenvalue equation

$$(4.22) \quad -\frac{1}{32} \frac{n^2}{n-1} \ddot{w} - \mu t^{2-\frac{4}{n}} w - nt^2 \Lambda w = 0,$$

where Λ is the eigenvalue.

This eigenvalue problem we also considered in a previous paper and proved that it has countably many solutions (w_i, Λ_i) with finite energy, i.e.,

$$(4.23) \quad \int_0^\infty \{ |\dot{w}_i|^2 + (1+t^2 + \mu t^{2-\frac{4}{n}}) |w_i|^2 \} < \infty,$$

cf. [2, Theorem 6.7].

We can then extend the spectral resolution which we proved in [3, Theorem 1.7] for a compact Cauchy hypersurface \mathcal{S}_0 to the case when \mathcal{S}_0 is non-compact:

4.3. Theorem. *Assume $n \geq 2$ and let \mathcal{S}_0 and the elliptic differential operator A satisfy the assumptions of the Theorem 4.2. Pick any solution (f, μ) of the eigenvalue problem (3.1), then there exist countably many solutions (w_i, Λ_i) of the implicit eigenvalue problem (4.22) such that*

$$(4.24) \quad \Lambda_i < \Lambda_{i+1} < \cdots < 0,$$

$$(4.25) \quad \lim_i \Lambda_i = 0,$$

and such that the functions

$$(4.26) \quad u_i = w_i f$$

are solutions of the wave equations (1.1) on page 1. The transformed eigenfunctions

$$(4.27) \quad \tilde{w}_i(t) = w_i(\lambda_i^{\frac{n}{4(n-1)}} t),$$

where

$$(4.28) \quad \lambda_i = (-\Lambda_i)^{-\frac{n-1}{n}},$$

form a basis of $L^2(\mathbb{R}_+, \mathbb{C})$ and also of the Hilbert space H defined as the completion of $C_c^\infty(\mathbb{R}_+, \mathbb{C})$ under the norm of the scalar product

$$(4.29) \quad \langle w, \tilde{w} \rangle_1 = \int_0^\infty \{\tilde{w}' \tilde{w}' + t^2 \tilde{w} \tilde{w}\},$$

where a prime or a dot denotes differentiation with respect to t .

4.4. Remark. This result is the best we can achieve under the present assumptions. In order to prove a mass gap, i.e., prove an estimate of the form

$$(4.30) \quad 0 < \epsilon_0 \leq \mu$$

for all eigenvalues μ of the eigenvalue equation (3.1) on page 8 we have to strengthen our assumptions on the zero order terms: Instead of the assumption (4.3) we have to require

$$(4.31) \quad -\frac{n}{2}R + G + \alpha_2 \frac{n}{2}m_0V \geq \epsilon_0 > 0 \quad \forall x \notin B_{R_1}(x_0),$$

then we immediately would derive a mass gap

An even stronger estimate of the form

$$(4.32) \quad -\frac{n}{2}R + G + \alpha_2 \frac{n}{2}m_0V \geq \epsilon_0 r^\delta \quad \forall x \notin B_{R_1}(x_0),$$

with $\delta > 0$, would yield that the operator A would have a pure point spectrum since the quadratic form

$$(4.33) \quad \langle Au, u \rangle$$

would then be compact relative to the L^2 -scalar product and we would be in the same situation as if \mathcal{S}_0 would be compact.

REFERENCES

- [1] I. M. Gel'fand and N. Ya. Vilenkin, *Generalized functions. Vol. 4: Applications of harmonic analysis*, Translated by Amiel Feinstein, Academic Press, New York - London, 1964.
- [2] Claus Gerhardt, *A unified field theory I: The quantization of gravity*, (2015), [arXiv:1501.01205](https://arxiv.org/abs/1501.01205).
- [3] ———, *A unified field theory II: Gravity interacting with a Yang-Mills and Higgs field*, (2016), [arXiv:1602.07191](https://arxiv.org/abs/1602.07191).
- [4] J.-L. Lions and E. Magenes, *Non-homogeneous boundary value problems and applications. Vol. I*, Springer-Verlag, New York, 1972, Translated from the French by P. Kenneth, Die Grundlehren der mathematischen Wissenschaften, Band 181.
- [5] Krzysztof Maurin, *Allgemeine Eigenfunktionsentwicklungen, unitäre Darstellungen lokalkompakter Gruppen und automorphe Funktionen*, Math. Ann. **165** (1966), 204–222.
- [6] ———, *Methods of Hilbert spaces*, Translated from the Polish by Andrzej Alexiewicz and Waclaw Zawadowski. Monografie Matematyczne, Tom 45, Państwowe Wydawnictwo Naukowe, Warsaw, 1967.
- [7] Michael Reed and Barry Simon, *Methods of modern mathematical physics. I. Functional analysis*, Academic Press, New York, 1972.
- [8] Helmut H. Schaefer, *Topological vector spaces*, The Macmillan Co., New York; Collier-Macmillan Ltd., London, 1966.

RUPRECHT-KARLS-UNIVERSITÄT, INSTITUT FÜR ANGEWANDTE MATHEMATIK, IM NEUENHEIMER FELD 294, 69120 HEIDELBERG, GERMANY

E-mail address: gerhardt@math.uni-heidelberg.de

URL: <http://www.math.uni-heidelberg.de/studinfo/gerhardt/>